

**DRAFT New Jersey
Greenhouse Gas Inventory and
Reference Case Projections 1990-2020**

February 2008

This document is presented by the Department of Environmental Protection in draft form and is being made available for public comment. Comments will be accepted and considered prior to its finalization.

Comments may be sent by March 20 to:
njglobalwarming@dep.state.nj.us

Executive Summary

This report was prepared in response to directives of Governor Corzine's Executive Order 54 and the New Jersey Global Warming Response Act. During the analysis conducted for this report, the Department of Environmental Protection (DEP) has benefitted from the extensive climate change research performed by Center for Climate Strategies (CCS), a 501c3 nonprofit organization. In addition, DEP conferred with the Center for Energy, Economic and Environmental Policy at Rutgers University and sought input from stakeholders and other state agencies, particularly the Board of Public Utilities (BPU), in developing these data.

This report presents a preliminary assessment of the State's greenhouse gas (GHG) emissions from 1990 to 2020. The inventory and forecast estimates serve as a starting point to assist the State with an initial comprehensive understanding of New Jersey's current and possible future GHG emissions, and thereby inform the identification and analysis of policy options for mitigating GHG emissions. The Global Warming Response Act directs DEP to evaluate and recommend such policy options later this year.

Emissions and Reference Case Projections (Business-as-Usual)

New Jersey's anthropogenic GHG emissions and anthropogenic sinks (carbon storage) were estimated for the period from 1990 to 2020. Historical GHG emission estimates (1990 through 2005) were developed using a set of generally accepted principles and guidelines for state GHG emissions estimates (both historical and forecasted), with adjustments as needed to provide New Jersey-specific data and inputs when it was possible to do so. The initial reference case projections (2006-2020) are based on a compilation of various existing projections of electricity generation, fuel use, and other GHG-emitting activities, along with a set of transparent assumptions.

Activities in New Jersey accounted for approximately 145.1 million metric tons (MMt) of *gross*¹ carbon dioxide equivalent (CO₂e) emissions (consumption basis) in 2005, an amount equal to about 2.0% of total US gross GHG emissions (based on 2005 US data).² New Jersey's gross GHG emissions are rising at a slower rate than those of the nation as a whole (gross emissions exclude carbon sinks, such as forests). New Jersey's gross GHG emissions increased by about 13% from 1990 to 2005, while national emissions rose by 16% from 1990 to 2005. The growth in New Jersey's emissions from 1990 to 2005 is primarily associated with the transportation and electricity consumption sectors.

Estimates of carbon sinks within New Jersey's forests have also been included in this report. For forests, the current estimates indicate that about 6.73 MMtCO₂e was stored in New Jersey forest biomass in 2005. This leads to *net* emissions of 138.3 MMtCO₂e in New Jersey in 2005. Figure ES-1 illustrates the State's emissions per capita and per unit of economic output. On a per capita basis, New Jersey residents emitted about 16.5 metric tons (Mt) of gross CO₂e in 1990, lower than the national average of 25.0 MtCO₂e in 1990. Per capita emissions in New Jersey increased slightly to 16.7 MtCO₂e/yr by 2005, while the per capita emissions for the US have

¹ Excluding GHG emissions removed due to forestry and other land uses.

² The national emissions used for these comparisons are based on 2005 emissions; (<http://www.epa.gov/climatechange/emissions/usinventoryreport.html>).

decreased slightly to 24.5 MtCO₂e/yr. As with the nation as a whole, economic growth exceeded emissions growth throughout the 1990-2005 period (leading to declining estimates of GHG emissions per unit of state product). From 1990 to 2005, emissions per unit of gross product dropped by 27% nationally, and by 22% in New Jersey.³

There are three principal sources of GHG emissions in New Jersey: transportation; electricity consumption; and the residential, commercial, and industrial (RCI) fuel use sectors. Transportation accounted for 34% of New Jersey's gross GHG emissions in 2005, while RCI fuel use accounted for 32% and electricity consumption contributed 24% of gross GHG emissions in 2005.

As illustrated in Figure ES-2 and shown numerically in Table ES-1, under the reference case projections, New Jersey's gross GHG emissions continue to grow, and are projected to climb to about 164.3 MMtCO₂e by 2020, reaching 28% above 1990 levels. As shown in Figure ES-3, the transportation sector is projected to be the largest contributor to future emissions growth in New Jersey, followed by electricity consumption and substitutes for ozone-depleting substances (ODS).

Some data gaps exist in this analysis, particularly for the reference case projections. Key refinements include review and revision of key emissions drivers that will be major determinants of New Jersey's future GHG emissions (such as the growth rate assumptions for electricity generation and consumption, transportation fuel use, and RCI fuel use). Appendices A through H provide the detailed methods, data sources, and assumptions for each GHG sector. Also included are descriptions of significant uncertainties in emission estimates or methods and suggested next steps for refinement of the inventory. Appendix I provides background information on GHGs and climate-forcing aerosols.

Potential Reductions

Throughout this report, potential reductions in emissions are noted where it is anticipated that such reductions can be achieved with strategies consistent with Governor Corzine's overall goal of reducing New Jersey's energy use by 20% below business as usual by 2020.⁴ These strategies include, but are not necessarily limited to, the Regional Greenhouse Gas Initiative (RGGI), New Jersey's Renewable Portfolio Standard (RPS), and the California Low Emissions Vehicle Program as adopted by New Jersey. Although not factored into the estimates of potential reductions recently enacted federal legislation⁵ will also have an effect on greenhouse gas emissions in the state. This category of "potential reductions is provided in this draft to give the reader a general understanding of how Governor Corzine's goals relate to current and projected greenhouse gas emissions. Elaboration on policies to achieve the Governor's goals as well as the

³ Based on real gross domestic product (millions of chained 2000 dollars) that excludes the effects of inflation, available from the US Bureau of Economic Analysis (<http://www.bea.gov/regional/gsp/>). The national emissions used for these comparisons are based on 2005 emissions. (<http://www.epa.gov/climatechange/emissions/usinventoryreport.html>).

⁴ See <http://www.nj.gov/emp/about/goals.html>.

⁵ The federal Energy Policy Act of 2007 requires refiners to use 9 billion gallons of grain ethanol in 2008 and 15 billion gallons annually by 2015. There also are separate new mandates for usage of biodiesel and fuels made from nonfood sources, such as crop waste and trees. The law also requires that by 2022 total biofuels use must reach 36 billion gallons annually.

methodology used to estimate their impact on greenhouse gas reductions will be provided in the near future as recommendations under the Energy Master Plan and Global Warming Response Act are presented for stakeholder input.

Table ES-1. New Jersey Historical and Reference Case GHG Emissions, by Sector^a

(Million Metric Tons CO ₂ e)	1990	2000	2005	2010	2020	Explanatory Notes for Projections
Energy	111.5	118.0	132.7	138.9	148.1	
Electricity, Production-Based	12.4	20.2	20.3	14.6	22.8	
Coal	6.93	10.67	9.59	7.42	8.85	See electric sector assumptions in Appendix A.
Natural Gas	3.64	7.41	8.32	5.49	12.4	
Oil	1.70	0.85	1.14	0.42	0.29	
Wood (CH ₄ and N ₂ O)	0.01	0.03	0.00	0.00	0.00	
MSW/LFG	0.17	1.27	1.27	1.27	1.27	
Net Imported Electricity	14.1	7.33	14.8	24.5	20.3	
Electricity Consumption Based	26.5	27.6	35.1	39.1	43.1	
Residential/Commercial/Industrial (RCI)	46.3	43.3	46.0	45.0	44.2	
Coal	0.70	0.033	0.029	0.029	0.030	Based on USDOE regional projections
Natural Gas	20.5	25.6	26.4	26.5	27.4	Based on NJ BPU projections
Oil	25.0	17.7	19.6	18.5	16.7	Based on NJ BPU projections
Wood (CH ₄ and N ₂ O)	0.13	0.08	0.07	0.06	0.06	Based on USDOE regional projections
Transportation	36.6	45.2	49.5	52.7	58.6	
On-road Gasoline	29.8	35.6	38.9	41.1	44.3	Based on USDOE regional projections
On-road Diesel	4.22	6.76	7.63	8.54	11.0	Based on USDOE regional projections
Marine Vessels	1.01	1.35	1.48	1.56	1.79	
Rail, Natural Gas, LPG, other	0.63	0.48	0.48	0.51	0.55	Based on USDOE regional projections
Jet Fuel and Aviation Gasoline	1.00	1.00	1.00	1.00	1.00	Estimated portion of emissions able to be controlled by NJ
Fossil Fuel Industry	2.06	1.87	2.02	2.06	2.14	
Natural Gas Industry	2.06	1.87	2.02	2.06	2.14	
Industrial Processes	1.26	2.86	4.02	5.49	8.61	
Nitric Acid Production (N ₂ O)	0.203	0.001	0.001	0.001	0.001	Based on State's modeling forecast of manufacturing employment for 2006-2020
Limestone and Dolomite Use (CO ₂)	0.000	0.003	0.005	0.005	0.004	Based on State's modeling forecast of manufacturing employment for 2006-2020
Soda Ash (CO ₂)	0.08	0.08	0.08	0.08	0.08	Based on 2004 and 2009 projections for U.S. production
ODS Substitutes (HFC, PFC)	0.010	2.41	3.59	5.16	8.37	EPA 2004 ODS cost study report
Electric Power T & D (SF ₆)	0.63	0.31	0.30	0.21	0.12	Based on national projections (USEPA)
Semiconductor Manufacturing (HFC, PFC, SF ₆)	0.01	0.03	0.03	0.02	0.01	Based on national projections (USEPA)
Laboratory Use of SF ₆	0.33	0.02	0.02	0.02	0.02	Assumed no change from 2005 levels.
Waste Management	13.4	6.67	6.76	6.46	6.01	
Waste Combustion	0.00	0.00	0.00	0.00	0.00	Captured under electricity production sector
Landfills	12.9	6.15	6.23	5.89	5.37	Includes waste landfilled out of state
Wastewater Management	0.45	0.52	0.54	0.57	0.64	Projections based on historical 1990 to 2005 average annual growth rate.
Agriculture	0.61	0.55	0.50	0.48	0.43	
Enteric Fermentation	0.13	0.09	0.08	0.07	0.06	
Manure Management	0.04	0.03	0.03	0.03	0.02	
Agricultural Soils	0.45	0.43	0.39	0.39	0.35	
Forestry and Land Use (Land Clearing Releases)	1.10	1.10	1.10	1.10	1.10	Based on NJDEP methodology; See Appendix H
Total Gross Emissions	127.8	129.1	145.1	152.5	164.3	
<i>increase relative to 1990</i>		<i>1%</i>	<i>13%</i>	<i>19%</i>	<i>28%</i>	
Forestry and Land Use (Sequestration)	-7.53	-6.99	-6.73	-6.46	-5.93	Based on NJDEP methodology; See Appendix H
Net Emissions (incl. forestry*)	120.3	122.1	138.3	146.0	158.3	

^aTotals may not equal exact sum of subtotals shown in this table due to independent rounding.

Figure ES-1. Historical New Jersey and US Gross GHG Emissions, Per Capita and Per Unit Gross Product






Figure ES-2. New Jersey Gross GHG Emissions by Sector, 1990-2020: Historical and Projected



RCI – direct fuel use in residential, commercial, and industrial sectors. ODS – ozone depleting substance.

Figure ES-3. Sector Contributions to Gross Emissions Growth in New Jersey, 1990-2020: Historic and Reference Case Projections (MMtCO₂e Basis)



RCI – direct fuel use in residential, commercial, and industrial sectors. ODS – ozone depleting substance. HFCs – hydrofluorocarbons.

Table ES-2. New Jersey GHG Emissions; Estimates and Projections (MMtCO₂e)^a

Sector	Sub-sector	2004	2020 BAU	2020 with potential reductions	Comments
Transportation	On-road gasoline	38.3	44.3	31.3	Projections sensitive to vehicle miles traveled (VMT) growth
	On-road diesel	7.5	11.0	10.5	Projections sensitive to VMT growth
	Aviation	1.0	1.0	1.0	Primarily jet fuel, estimated in-state use; total is about 8 metric tons
	Marine	1.5	1.8	1.8	Diesel and residual; does not include port expansion
	Railroad & Other	0.5	0.6	0.6	
Electricity Generation	In-state	19.0	21.5	19.8	2020 with actions set equal to RGGI CO ₂ emissions cap
	In-state, from MSW	1.3	1.3	1.3	
	Imported	13.4	20.3	4.8	2020 with actions is minus out-of-state renewables from RPS
Residential	Space heat	14.2	15.5	12.4	
	Other combustion	3.4	3.8	3.0	
Commercial	Space heat	6.1	5.7	4.6	
	Other combustion	4.8	3.6	2.9	
Industrial	Space heat	0.7	0.4	0.3	
	Other combustion	17.1	15.3	13.9	
Halogenated gases (excluding SF ₆)		3.4	8.4	8.4	
SF ₆		0.3	0.1	0.1	
Industrial non-fuel related		0.1	0.1	0.1	
Agriculture		0.5	0.4	0.4	
Natural gas T&D		2.0	2.1	2.1	
Landfills, POTWs		6.8	6.0	6.0	Includes emissions from out-of-state landfills taking NJ MSW
Released through land clearing		1.1	1.1	1.1	
Total Gross Emissions		142	164	125	
Sequestered by forests		-6.8	-5.9	-5.9	
Net Emissions (including forests)		136	158	120	

NOTE: Values shown are estimates only, and are subject to revision as better data become available. 2004 numbers are believed accurate to within 5%; there is considerably more uncertainty with 2020 projections.

^a Totals may not equal exact sum of subtotals shown in this table due to independent rounding. BAU – Business-as-Usual, MSW - municipal solid waste, POTW - Publicly Owned Treatment Works, RPS - Renewable Portfolio Standard, RGGI - Regional Greenhouse Gas Initiative, SF₆ – sulfur hexafluoride, T&D – transmission and distribution.

^b There are no specific state RGGI caps. Projected number shown is based on NJ's portion of the total regional cap. Actual in-state emissions may differ. Cap only applies to the approximately 95% of non-MSW generation capacity that is subject to RGGI.

Table of Contents

Executive Summary	iii
Acronyms and Key Terms	6
Summary of Preliminary Findings	10
Introduction	10
New Jersey Greenhouse Gas Emissions: Sources and Trends	11
Historical Emissions	13
Overview	13
A Closer Look at the Three Major Sources: Transportation, RCI Fuel Use Sector, and Electricity Supply	15
Reference Case Projections (Business as Usual)	16
Potential Reductions	17
Key Uncertainties and Next Steps	19
Approach	20
General Methodology	20
General Principles and Guidelines	21
Appendix A. Energy Supply	A-1
Appendix B. Residential, Commercial, and Industrial (RCI) Fuel Combustion	B-1
Appendix C. Transportation Energy Use	C-1
Appendix D. Industrial Processes	D-1
Appendix E. Fossil Fuel Extraction and Distribution Industry	E-1
Appendix F. Agriculture	F-1
Appendix G. Waste Management	G-1
Appendix H. Forestry and Land-Use Change	H-1

Acronyms and Key Terms

AEO2006 – EIA’s Annual Energy Outlook 2006

Ag – Agriculture

BAU – Business as Usual

bbls – Barrels

BC – Black Carbon*

Bcf – Billion cubic feet

BOD – Biochemical Oxygen Demand

BTU – British thermal unit

C – Carbon

CaCO₃ – Calcium Carbonate

CCS – Center for Climate Strategies

CEC – Commission for Environmental Cooperation

CEEEP – Center for Energy, Economic & Environmental Policy (Rutgers University)

CFCs – Chlorofluorocarbons

CH₄ – Methane

CO – Carbon Monoxide

CO₂ – Carbon Dioxide

CO₂e – Carbon Dioxide equivalent

COLE – Carbon Online Estimator

CRP – Federal Conservation Reserve Program

CRSSA – Center for Remote Sensing and Spatial Analysis

EC – Elemental Carbon

EEZ – Exclusive Economic Zone

EIA – US DOE Energy Information Administration
EIIP – Emissions Inventory Improvement Program
EMP – Energy Master Plan
FAA – Federal Aviation Administration
FAPRI – Food and Agricultural Policy Research Institute
FHWA – Federal Highway Administration
FIA – Forest Inventory Analysis
GHG – Greenhouse Gases
GIS – Geographical Information System
GWh – Gigawatt-hour
GWP – Global Warming Potential
HFCs – Hydrofluorocarbons
IPCC – Intergovernmental Panel on Climate Change
kWh – kilowatt-hour
LandGEM – Landfill Gas Emissions Model
lb – pound
LF – Landfill
LFGTE – Landfill Gas Collection System and Landfill-Gas-to-Energy
LNG – Liquefied Natural Gas
LPG – Liquefied Petroleum Gas
MMBtu – Million British thermal units
MMt – Million Metric tons
MMtCO₂e – Million Metric tons Carbon Dioxide equivalent
Mt – Metric ton (equivalent to 1.102 short tons)

MSW – Municipal Solid Waste

MW – Megawatt

MWh – Megawatt-hour

N – Nitrogen

N₂O – Nitrous Oxide

NEI – National Emissions Inventory

NO₂ – Nitrogen Dioxide

NO_x – Nitrogen Oxides

NASS – National Agricultural Statistics Service

NJDEP – New Jersey Department of Environmental Protection

NJDOT – New Jersey Department of Transportation

NMVOCs – Nonmethane Volatile Organic Compounds

O₃ – Ozone

ODS – Ozone-Depleting Substances

OM – Organic Matter

PFCs – Perfluorocarbons

PM – Particulate Matter

POTW – Publicly Owned Treatment Works

ppb – parts per billion

ppm – parts per million

ppt – parts per trillion

RCI – Residential, Commercial, and Industrial

RGGI – Regional Greenhouse Gas Initiative

RPS – Renewable Portfolio Standard

SAR – Second Assessment Report

SED – State Energy Data

SF₆ – Sulfur Hexafluoride

SGIT – State Greenhouse Gas Inventory Tool

Sinks – Removals of carbon from the atmosphere, with the carbon stored in forests, soils, landfills, wood structures, or other biomass-related products.

TAR – Third Assessment Report

T&D – Transmission and Distribution

TBtu – Trillion British Thermal Units

TOG – Total Organic Carbon

UNFCCC – United Nations Framework Convention on Climate Change

US EPA – United States Environmental Protection Agency

US DOE – United States Department of Energy

USDA – United States Department of Agriculture

USFS – United States Forest Service

USGS – United States Geological Survey

VMT – Vehicle-Miles Traveled

W/m² – Watts per Square Meter

WMO – World Meteorological Organization

WW – Wastewater

yr – year

Summary of Preliminary Findings

Introduction

This report was prepared in response to directives of Governor Corzine’s Executive Order 54 and the New Jersey Global Warming Response Act. During the analysis conducted for this report, the Department of Environmental Protection (DEP) has benefitted from the extensive climate change research performed by the Center for Climate Strategies (CCS), a 501c3 nonprofit organization. In addition, DEP conferred with the Center for Energy, Economic and Environmental Policy at Rutgers University and sought input from stakeholders and other state agencies in developing these data.

This report presents a preliminary assessment of the State’s greenhouse gas (GHG) emissions from 1990 to 2020. The inventory and forecast estimates serve as a starting point to assist the State with an initial comprehensive understanding of New Jersey’s current and possible future GHG emissions, and thereby inform the identification and analysis of policy options for mitigating GHG emissions. The Global Warming Response Act directs DEP to evaluate and recommend such policy options later this year.

New Jersey’s anthropogenic GHG emissions and anthropogenic sinks (carbon storage) were estimated for the period from 1990 to 2020. Historical GHG emission estimates (1990 through 2005)⁶ were developed using a set of generally accepted principles and guidelines for State GHG emissions inventories, as described in the “Approach” section below, relying to the extent possible on New Jersey-specific data and inputs. The initial reference case projections (2006-2020) are based on a compilation of various existing projections of electricity generation, fuel use, and other GHG-emitting activities, along with a set of simple, transparent assumptions described in the appendices of this report.

This report covers the six gases included in the US Greenhouse Gas Inventory: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulfur hexafluoride (SF₆). Emissions of these GHGs are presented using a common metric, CO₂ equivalence (CO₂e), which indicates the relative contribution of each gas to global average radiative forcing on a Global Warming Potential- (GWP) weighted basis.

It is important to note that the preliminary emissions estimates reflect the *GHG emissions associated with the electricity sources used to meet New Jersey’s demands*, corresponding to a consumption-based approach to emissions accounting (see “Approach” section below). Another way to look at electricity emissions is to consider the *GHG emissions produced by electricity generation facilities in the State*. This report covers both methods of accounting for emissions, but for consistency, all total results are reported as *consumption-based*.

⁶ The last year of available historical data varies by sector; ranging from 2000 to 2006.

New Jersey Greenhouse Gas Emissions: Sources and Trends

Table 1 provides a summary of GHG emissions estimated for New Jersey by sector for the years 1990, 2000, 2005, 2010, and 2020. Details on the methods and data sources used to construct these draft estimates are provided in the appendices to this report. In the sections below, we discuss GHG emission sources (positive, or *gross*, emissions) and sinks (negative emissions) separately in order to identify trends, projections, and uncertainties clearly for each.

This next section of the report provides a summary of the historical emissions (1990 through 2005) followed by a summary of the reference-case projection-year emissions (2006 through 2020) and key uncertainties. We also provide an overview of the general methodology, principles, and guidelines followed for preparing the inventories. Appendices A through H provide the detailed methods, data sources, and assumptions for each GHG sector.

Table 1. New Jersey Historical and Reference Case GHG Emissions, by Sector^a

(Million Metric Tons CO ₂ e)	1990	2000	2005	2010	2020	Explanatory Notes for Projections
Energy	111.5	118.0	132.7	138.9	148.1	
Electricity, Production-Based	12.4	20.2	20.3	14.6	22.8	
Coal	6.93	10.67	9.59	7.42	8.85	See electric sector assumptions in Appendix A.
Natural Gas	3.64	7.41	8.32	5.49	12.4	
Oil	1.70	0.85	1.14	0.42	0.29	
Wood (CH ₄ and N ₂ O)	0.01	0.03	0.00	0.00	0.00	
MSW/LFG	0.17	1.27	1.27	1.27	1.27	
Net Imported Electricity	14.1	7.33	14.8	24.5	20.3	
Electricity Consumption Based	26.5	27.6	35.1	39.1	43.1	
Residential/Commercial/Industrial (RCI)	46.3	43.3	46.0	45.0	44.2	
Coal	0.70	0.033	0.029	0.029	0.030	Based on USDOE regional projections
Natural Gas	20.5	25.6	26.4	26.5	27.4	Based on NJ BPU projections
Oil	25.0	17.7	19.6	18.5	16.7	Based on NJ BPU projections
Wood (CH ₄ and N ₂ O)	0.13	0.08	0.07	0.06	0.06	Based on USDOE regional projections
Transportation	36.6	45.2	49.5	52.7	58.6	
On-road Gasoline	29.8	35.6	38.9	41.1	44.3	Based on USDOE regional projections
On-road Diesel	4.22	6.76	7.63	8.54	11.0	Based on USDOE regional projections
Marine Vessels	1.01	1.35	1.48	1.56	1.79	
Rail, Natural Gas, LPG, other	0.63	0.48	0.48	0.51	0.55	Based on USDOE regional projections
Jet Fuel and Aviation Gasoline	1.00	1.00	1.00	1.00	1.00	Estimated portion of emissions able to be controlled by NJ
Fossil Fuel Industry	2.06	1.87	2.02	2.06	2.14	
Natural Gas Industry	2.06	1.87	2.02	2.06	2.14	
Industrial Processes	1.26	2.86	4.02	5.49	8.61	
Nitric Acid Production (N ₂ O)	0.203	0.001	0.001	0.001	0.001	Based on State's modeling forecast of manufacturing employment for 2006-2020
Limestone and Dolomite Use (CO ₂)	0.000	0.003	0.005	0.005	0.004	Based on State's modeling forecast of manufacturing employment for 2006-2020
Soda Ash (CO ₂)	0.08	0.08	0.08	0.08	0.08	Based on 2004 and 2009 projections for U.S. production
ODS Substitutes (HFC, PFC)	0.010	2.41	3.59	5.16	8.37	EPA 2004 ODS cost study report
Electric Power T & D (SF ₆)	0.63	0.31	0.30	0.21	0.12	Based on national projections (USEPA)
Semiconductor Manufacturing (HFC, PFC, SF ₆)	0.01	0.03	0.03	0.02	0.01	Based on national projections (USEPA)
Laboratory Use of SF ₆	0.33	0.02	0.02	0.02	0.02	Assumed no change from 2005 levels.
Waste Management	13.4	6.67	6.76	6.46	6.01	
Waste Combustion	0.00	0.00	0.00	0.00	0.00	Captured under electricity production sector
Landfills	12.9	6.15	6.23	5.89	5.37	Includes waste landfilled out of state
Wastewater Management	0.45	0.52	0.54	0.57	0.64	Projections based on historical 1990 to 2005 average annual growth rate.
Agriculture	0.61	0.55	0.50	0.48	0.43	
Enteric Fermentation	0.13	0.09	0.08	0.07	0.06	
Manure Management	0.04	0.03	0.03	0.03	0.02	
Agricultural Soils	0.45	0.43	0.39	0.39	0.35	
Forestry and Land Use (Land Clearing Releases)	1.10	1.10	1.10	1.10	1.10	Based on NJDEP methodology; See Appendix H
Total Gross Emissions	127.8	129.1	145.1	152.5	164.3	
<i>increase relative to 1990</i>		<i>1%</i>	<i>13%</i>	<i>198%</i>	<i>28%</i>	
Forestry and Land Use (Sequestration)	-7.53	-6.99	-6.73	-6.46	-5.93	Based on NJDEP methodology; See Appendix H
Net Emissions (incl. forestry*)	120.3	122.1	138.3	146.0	158.3	

^aTotals may not equal exact sum of subtotals shown in this table due to independent rounding.


Historical Emissions

Overview

Preliminary analyses suggest that in 2005, activities in New Jersey accounted for approximately 145.1 million metric tons (MMt) of gross CO₂e emissions (consumption basis), an amount equal to about 2.0% of total US gross GHG emissions (based on 2005 US data).⁷ New Jersey's gross GHG emissions are rising at a slower rate than those of the nation as a whole (gross emissions exclude carbon sinks, such as forests). New Jersey's gross GHG emissions increased by about 13% from 1990 to 2005, while national emissions rose by 16% from 1990 to 2005.

Figure 1 illustrates the State's emissions per capita and per unit of economic output. On a per capita basis, New Jersey residents emitted about 16.5 metric tons (Mt) of CO₂e in 1990, lower than the national average of 25.0 MtCO₂e in 1990. Per capita emissions in New Jersey increased slightly to 16.7 MtCO₂e/yr by 2005, while the per capita emissions for the US have decreased slightly to 24.5 MtCO₂e/yr. As with the nation as a whole, economic growth exceeded emissions growth throughout the 1990-2005 period (leading to declining estimates of GHG emissions per unit of state product). From 1990 to 2005, emissions per unit of gross product dropped by 27% nationally, and by 22% in New Jersey.⁸

Figure 1. Historical New Jersey and US Gross GHG Emissions, Per Capita and Per Unit Gross Product




⁷ United States emissions estimates are drawn from US EPA 2007, *Inventory of US Greenhouse Gas Emissions and Sinks: 1990 to 2005*; (<http://www.epa.gov/climatechange/emissions/usinventoryreport.html>).

⁸ Based on real gross domestic product (millions of chained 2000 dollars) that excludes the effects of inflation, available from the US Bureau of Economic Analysis (<http://www.bea.gov/regional/gsp/>). The national emissions used for these comparisons are based on 2005 emissions. (<http://www.epa.gov/climatechange/emissions/usinventoryreport.html>).

Figure 2 compares gross GHG emissions estimated for New Jersey to emissions for the US for year 2000. Principal sources of New Jersey's GHG emissions are transportation and the residential, commercial, and industrial (RCI) fuel use sectors, accounting for 35% and 34% of New Jersey's gross GHG emissions in 2000, respectively. The next largest contributor is electricity consumption, accounting for 21% of gross GHG emissions in 2000. The waste management and agriculture sectors each contribute 5.2% and 0.4% of gross GHG emissions in 2000, respectively.

Industrial process emissions comprised 2.2% of State GHG emissions in 2000. Although industrial process emissions are rising rapidly due to the increasing use of HFC as substitutes for ozone-depleting chlorofluorocarbons (CFCs), their overall contribution is estimated to be only 5.2% of New Jersey's gross GHG emissions in 2020 due to significant growth in other sectors.⁹ Other industrial process emissions result from CO₂ released during soda ash, limestone, and dolomite use; releases of SF₆ from transformers used in the transmission and distribution of electricity and laboratory uses; and N₂O emissions associated with the manufacture of nitric acid. Methane emissions associated with natural gas transmission and distribution (T&D) (included under the fossil fuel industry category) accounted for 1.4% of the State's gross GHG emissions in 2000. The forestry sector accounts for 0.9% of New Jersey's gross GHG emissions in 2000. These are emissions that occur when forested and vegetated lands are cleared for other purposes, such as the construction of buildings and parking lots.

Figure 2. Gross GHG Emissions by Sector, 2000, New Jersey and US



NOTE: at a national level, forests act as a net sink of CO₂; therefore, they do not show up in the above graph of gross U.S. emissions sources.

⁹ CFCs are also potent GHGs; they are not, however, included in GHG estimates because of concerns related to implementation of the Montreal Protocol (See Appendix I for additional information). HFCs are used as refrigerants in the RCI and transport sectors as well as in the industrial sector; they are included here, however, within the industrial processes emissions.

Overall, however, forestry activities in New Jersey are estimated to be net sinks of GHG emissions throughout the period from 1990 through 2020. Through sequestration, forested lands in New Jersey are expected to store 6.7 MMtCO_{2e} in 2005, decreasing to 5.9 MMtCO_{2e} by 2020.

A Closer Look at the Three Major Sources: Transportation, RCI Fuel Use Sector, and Electricity Supply

Transportation Sector

As shown in Figure 2, the transportation sector accounted for about 35% of New Jersey's gross GHG emissions in 2000 (about 45.2 MMtCO_{2e}), which was higher than the national average share of emissions from transportation fuel consumption (26%). The GHG emissions associated with New Jersey's transportation sector increased by 12.9 MMtCO_{2e} between 1990 and 2005, accounting for about 75% of the State's net growth in gross GHG emissions in this time period.

From 1990 through 2005, New Jersey's GHG emissions from transportation fuel use have risen steadily at an average rate of about 2.0% annually. In 2005, on-road gasoline vehicles accounted for about 79% of transportation GHG emissions. On-road diesel vehicles accounted for another 15% of emissions, and marine vessels for roughly 3%. Air travel, rail, and other sources (natural gas- and liquefied petroleum gas- (LPG-) fueled-vehicles used in transport applications) accounted for the remaining 3% of transportation emissions. As a result of New Jersey's population and economic growth and an increase in total vehicle miles traveled (VMT) during the 1990s, on-road gasoline use grew 34% between 1990 and 2005. Meanwhile, on-road diesel use rose 81% during that period, suggesting an even more rapid growth in freight movement within or across the State. Marine fuel use also increased by about 46% from 1990 to 2005. Rail fuel use decreased about 44% from 1990 to 2005.

Residential, Commercial, and Industrial (RCI) Sector

Activities in the RCI¹⁰ sectors produce GHG emissions when fuels are combusted to provide space heating, process heating, and other applications. In 2000, combustion of oil, natural gas, coal, and wood in the RCI sectors contributed about 34% of New Jersey's gross GHG emissions, much higher than RCI sector contribution for the nation (23%).

By 2005, the RCI sector emissions were at about 32% (46.0 MMtCO_{2e}) of gross GHG emissions. In 2005, the residential sector's share of total RCI emissions from direct fuel use was 38% (17.7 MMtCO_{2e}), the commercial sector accounted for 23% (10.7 MMtCO_{2e}), and the industrial sector's share of total RCI emissions from direct fuel use was 38% (17.6 MMtCO_{2e}). Overall, emissions for the RCI sectors (excluding those associated with electricity consumption) are expected to decrease by 3.9% between 2005 and 2020. Emissions from the residential sector are projected to increase by 9% between 2005 and 2020. In contrast, emissions from the commercial and industrial sectors are expected to decrease by 13% and 11%, respectively, between 2005 and 2020.

¹⁰ The industrial sector includes emissions associated with agricultural energy use and fuel used by the fossil fuel extraction and distribution industry.

Electricity Supply Sector

New Jersey is a net importer of electricity, meaning that the State consumes more electricity than is produced in the State. For this analysis, it was assumed that all power generated in New Jersey was consumed in New Jersey, and that remaining electricity demand was met by imported power. Imported power accounted for 28% of the electricity consumed in New Jersey in 2004.¹¹ Emissions from the power produced in-state are dominated by coal and natural gas fuel use. As shown in Figure 2, electricity consumption accounted for about 21% of New Jersey's gross GHG emissions in 2000 (about 27.6 MMtCO₂e), which was lower than the national average share of emissions from electricity consumption (32%).¹² The GHG emissions associated with New Jersey's electricity sector increased by 8.6 MMtCO₂e between 1990 and 2005, accounting for 50% of the state's net growth in gross GHG emissions in this time period.

In 2005, emissions associated with New Jersey's electricity consumption (35.1 MMtCO₂e) were about 14.8 MMtCO₂e higher than those associated with electricity production (20.3 MMtCO₂e). The higher level for consumption-based emissions reflects GHG emissions associated with net imports of electricity to meet New Jersey's electricity demand.¹³ Projections of electricity sales for 2005 through 2020 indicate that New Jersey will remain a net importer of electricity. For the period covering 2005 through 2020, the reference case projection assumes that production-based emissions (associated with electricity generated in-state) will increase by about 2.5 MMtCO₂e, and consumption-based emissions (associated with electricity consumed in-state) will increase by about 8.0 MMtCO₂e.

The consumption-based approach can better reflect the emissions (and emissions reductions) associated with activities occurring in New Jersey, particularly with respect to electricity use (and efficiency improvements), and is particularly useful for policy-making.

Reference Case Projections (Business as Usual)

Relying on a variety of sources for projections, as noted below and in the appendices, we developed a simple reference case projection of GHG emissions through 2020. As illustrated in Figure 3 and shown numerically in Table 1, under the reference case projections, New Jersey's gross GHG emissions continue to grow steadily, climbing to about 164.3 MMtCO₂e by 2020, 28% above 1990 levels. The transportation sector is projected to be the largest contributor to future emissions growth, followed by emissions associated with electricity consumption and then by ODS Substitutes (HFCs), as shown in Figure 4. Table 2 provides a sector-level summary of emissions for 2004 (the latest year for which activity data were available), 2020 business-as-usual, and 2020 with potential reductions.

¹¹ Based on EIA 2004 State Electricity Profiles; total NJ retail sales were 77,593,167 MWh and net NJ generation was 55,882,342 MWh in 2004.


¹² For the US as a whole, there is relatively little difference between the emissions from electricity use and emissions from electricity production, as the US imports only about 1% of its electricity, and exports far less. New Jersey's situation is different, since it is a net electricity importer.

¹³ Estimating the emissions associated with electricity use requires an understanding of the electricity sources (both in-state and out-of-state) used by utilities to meet consumer demand. The current estimate reflects some very simple assumptions, as described in Appendix A.

Potential Reductions

Throughout this report, potential reductions in emissions are noted where it is anticipated that such reductions can be achieved with strategies consistent with Governor Corzine’s overall goal of reducing New Jersey’s energy use by 20% below business as usual by 2020.¹⁴ These strategies include, but are not necessarily limited to, the Regional Greenhouse Gas Initiative (RGGI), New Jersey’s Renewable Portfolio Standard (RPS), and the California Low Emissions Vehicle Program as adopted by New Jersey. Although not factored into the estimates of potential reductions recently enacted federal legislation¹⁵ will also have an effect on greenhouse gas emissions in the state. . This category of “potential reductions is provided in this draft to give the reader a general understanding of how Governor Corzine’s goals relate to current and projected greenhouse gas emissions. Elaboration on policies to achieve the Governor’s goals as well as the methodology used to estimate their impact on greenhouse gas reductions will be provided in the near future as recommendations under the Energy Master Plan and Global Warming Response Act are presented for stakeholder input.

Figure 3. New Jersey Gross GHG Emissions by Sector, 1990-2020: Historical and Projected




RCI – direct fuel use in residential, commercial, and industrial sectors. ODS – ozone depleting substance.

¹⁴ See <http://www.nj.gov/emp/about/goals.html>.

¹⁵ The federal Energy Policy Act of 2007 requires refiners to use 9 billion gallons of grain ethanol in 2008 and 15 billion gallons annually by 2015. There also are separate new mandates for usage of biodiesel and fuels made from nonfood sources, such as crop waste and trees. The law also requires that by 2022 total biofuels use must reach 36 billion gallons annually.

Figure 4. Sector Contributions to Gross Emissions Growth in New Jersey, 1990-2020: Historic and Reference Case Projections (MMtCO₂e Basis)



RCI – direct fuel use in residential, commercial, and industrial sectors. ODS – ozone depleting substance. HFCs – hydrofluorocarbons.

Table 2. New Jersey GHG Emissions; Estimates and Projections (MMtCO₂e)^a

Sector	Sub-sector	2004	2020 BAU	2020 with potential reductions	Comments
Transportation	On-road gasoline	38.3	44.3	31.3	Projections sensitive to vehicle miles traveled (VMT) growth
	On-road diesel	7.5	11.0	10.5	Projections sensitive to VMT growth
	Aviation	1.0	1.0	1.0	Primarily jet fuel, estimated in-state use; total is about 8 metric tons
	Marine	1.5	1.8	1.8	Diesel and residual; does not include port expansion
	Railroad & Other	0.5	0.6	0.6	
Electricity Generation	In-state	19.0	21.5	19.8	2020 with actions set equal to RGGI CO ₂ emissions cap
	In-state, from MSW	1.3	1.3	1.3	
	Imported	13.4	20.3	4.8	2020 with actions is minus out-of-state renewables from RPS
Residential	Space heat	14.2	15.5	12.4	
	Other combustion	3.4	3.8	3.0	
Commercial	Space heat	6.1	5.7	4.6	
	Other combustion	4.8	3.6	2.9	
Industrial	Space heat	0.7	0.4	0.3	
	Other combustion	17.1	15.3	13.9	
Halogenated gases (excluding SF ₆)		3.4	8.4	8.4	
SF ₆		0.3	0.1	0.1	
Industrial non-fuel related		0.1	0.1	0.1	
Agriculture		0.5	0.4	0.4	
Natural gas T&D		2.0	2.1	2.1	
Landfills, POTWs		6.8	6.0	6.0	Includes emissions from out-of-state landfills taking NJ MSW
Released through land clearing		1.1	1.1	1.1	
Total Gross Emissions		142	164	125	
Sequestered by forests		-6.8	-5.9	-5.9	
Net Emissions (including forests)		136	158	120	

NOTE: Values shown are estimates only, and are subject to revision as better data become available. 2004 numbers are believed accurate to within 5%; there is considerably more uncertainty with 2020 projections.

^a Totals may not equal exact sum of subtotals shown in this table due to independent rounding. MSW - municipal solid waste, POTW - Publicly Owned Treatment Works, RPS - Renewable Portfolio Standard, RGGI - Regional Greenhouse Gas Initiative, SF₆ – sulfur hexafluoride, T&D – transmission and distribution.

^b There are no specific state RGGI caps. Projected number shown is based on NJ's portion of the total regional cap. Actual in-state emissions may differ. Cap only applies to the approximately 95% of non-MSW generation capacity that is subject to RGGI.

Key Uncertainties and Next Steps

Some data gaps exist in this inventory, and particularly in the reference case projections. Key tasks for future refinement of this inventory and forecast include review and revision of key drivers, such as the transportation, electricity demand, and RCI fuel use growth rates that will be major determinants of New Jersey's future GHG emissions (See Table 3 and Figure 4). These

growth rates are driven by uncertain economic, demographic and land use trends (including growth patterns and transportation system impacts), all of which deserve closer review and discussion.

Table 3. Key Annual Growth Rates for New Jersey, Historical and Projected

	1990-2005	2005-2020	Sources
Population^a	0.77%	0.68%	Center for Energy, Economic & Environmental Policy (CEEPP, 2007). ¹⁶
Employment^a Goods Services	NA ^b NA	-0.8% 1.2%	Historical employment annual growth rates not available. Annual growth rates for 2005-2020 based on Energy modeling for New Jersey (see reference for population forecast).
Electricity Sales Total Sales^c NJ Sales^d Imported Sales^e	1.8% 2.8% -0.5%	1.9% 1.5% 2.9%	Historical annual growth rates calculated using data for 1990 and 2005 from EIA's State Electricity Profiles for each year. Annual growth rates for 2005-2020 based on Energy modeling for New Jersey. ¹⁷
Vehicle Miles Traveled	1.5%	1.2%	Historical data from New Jersey Department of Transportation. Growth rates calculated based on linear regression of historical data.

^a For the RCI fuel consumption sectors, population and employment projections for New Jersey were used together with US DOE EIA's Annual Energy Outlook 2006 (AEO2006) projections of changes in fuel use for the EIA's Middle Atlantic region on a per capita basis for the residential sector, and on a per employee basis for the commercial and industrial sectors. For instance, growth in New Jersey's residential natural gas use is calculated as the New Jersey population growth times the change in per capita natural gas use for the Middle Atlantic region.

^b NA – Not available; historical employment data for New Jersey for the goods producing and services providing sectors could not be identified during development of this report.

^c Represents annual growth in total sales of electricity by generators in and outside of New Jersey to RCI sectors located within New Jersey (consumption basis).

^d Represents annual growth in total sales of electricity by generators in New Jersey to RCI sectors located within New Jersey (production basis). Annual growth rate calculated using data for 1990 and 2004.

^e Represents annual growth rate in sales of electricity imported into New Jersey.

Approach

The principal goal of compiling the inventories and reference case projections presented in this document is to provide NJDEP with a general understanding of New Jersey's historical, current, and projected (expected) GHG emissions. The following sections explain the general methodology and the general principles and guidelines followed during development of these GHG inventories for New Jersey.

General Methodology

The analyses contained in this report were developed in consultation with other state agencies, most notably, the Board of Public Utilities, and in conference with The Center for Climate Strategies, The Rutgers University, Center for Energy, Economic and Environmental Policy and stakeholders. The overall goal of this effort is to provide simple and straightforward estimates, with an emphasis on robustness, consistency, and transparency. As a result, we rely on reference

¹⁶ CEEPP, 2007, Center for Energy, Economic & Environmental Policy (CEEPP), Edward J. Bloustein School of Planning and Public Policy, Rutgers, The State University of New Jersey, 33 Livingston Ave., Room 154, New Brunswick, NJ 08901. Data provided by Nancy Mantell to CCS on March 30, 2007 (file named "Electricity Prelim Data (3-30-07).xls").

¹⁷ CEEPP, 2007, Data provided by Andrew Contrell to Michael Aucott, NJ DEP, September 2007.

forecasts from best available State and regional sources where possible. Where reliable existing forecasts are lacking, we use straightforward spreadsheet analysis and constant growth-rate extrapolations of historical trends rather than complex modeling.

In most cases, we follow the same approach to emissions accounting for historical inventories used by the US EPA in its national GHG emissions inventory¹⁸ and its guidelines for States.¹⁹ These inventory guidelines were developed based on the guidelines from the Intergovernmental Panel on Climate Change (IPCC), the international organization responsible for developing coordinated methods for national GHG inventories.²⁰ The inventory methods provide flexibility to account for local conditions. The key sources of activity and projection data used are shown in Table 4. Table 4 also provides the descriptions of the data provided by each source and the uses of each data set in this analysis.

General Principles and Guidelines

A key part of this effort involves the establishment and use of a set of generally accepted accounting principles for evaluation of historical and projected GHG emissions, as follows:

- **Transparency:** We report data sources, methods, and key assumptions to allow open review and opportunities for additional revisions later based on input from others. In addition, we will report key uncertainties where they exist.
- **Consistency:** To the extent possible, the inventory and projections will be designed to be externally consistent with current or likely future systems for State and national GHG emission reporting. We have used the EPA tools for State inventories and projections as a starting point. These initial estimates were then augmented and/or revised as needed to conform with State-based inventory and base-case projection needs. For consistency in making reference case projections, we define reference case actions for the purposes of projections as those *currently in place or reasonably expected over the time period of analysis*.
- **Priority of Existing State and Local Data Sources:** In gathering data and in cases where data sources conflicted, we placed highest priority on local and State data and analyses, followed by regional sources, with national data or simplified assumptions such as constant linear extrapolation of trends used as defaults where necessary.
- **Priority of Significant Emissions Sources:** In general, activities with relatively small emissions levels may not be reported with the same level of detail as other activities.
- **Comprehensive Coverage of Gases, Sectors, State Activities, and Time Periods.** This analysis aims to comprehensively cover GHG emissions associated with activities in New Jersey. It covers all six GHGs covered by US and other national inventories: CO₂, CH₄, N₂O, SF₆, HFCs, and PFCs. The inventory estimates are for the year 1990, with subsequent years included up to most recently available data (typically 2002 to 2005), with projections to 2010 and 2020.
- **Use of Consumption-Based Emissions Estimates:** To the extent possible, we estimated emissions that are caused by activities that occur in New Jersey. For example, we reported emissions associated with the electricity consumed in New Jersey. The rationale

¹⁸ US EPA 2007, *Inventory of US Greenhouse Gas Emissions and Sinks: 1990 to 2005*; (<http://www.epa.gov/climatechange/emissions/usinventoryreport.html>).

¹⁹ <http://yosemite.epa.gov/oar/globalwarming.nsf/content/EmissionsStateInventoryGuidance.html>.

²⁰ <http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.htm>.

for this method of reporting is that it can more accurately reflect the impact of State-based policy strategies such as energy efficiency on overall GHG emissions, and it resolves double-counting and exclusion problems with multi-emissions issues. This approach can differ from how inventories are compiled, for example, on an in-state production basis, in particular for electricity.

Table 4. Key Sources for New Jersey Data, Inventory Methods, and Growth Rates

Source	Information provided	Use of Information in this Analysis
US EPA State Greenhouse Gas Inventory Tool (SGIT)	US EPA SGIT is a collection of linked spreadsheets designed to help users develop State GHG inventories for 1990-2005. US EPA SGIT contains default data for each State for most of the information required for an inventory. The SGIT methods are based on the methods provided in the Volume VIII document series published by the Emissions Inventory Improvement Program (http://www.epa.gov/ttn/chief/eiip/techreport/volume08/index.html).	Where not indicated otherwise, SGIT is used to calculate emissions for 1990-2005 from RCI fuel combustion, transportation, industrial processes, agriculture and forestry, and waste. We use SGIT emission factors (CO ₂ , CH ₄ , and N ₂ O per BTU consumed) to calculate energy use emissions.
US DOE Energy Information Administration (EIA) State Energy Data (SED)	EIA SED provides energy use data in each State, annually to 2004 for all fuels, and to 2005 for natural gas and wood.	EIA SED is the source for most energy use data. Emission factors from US EPA SGIT are used to calculate energy-related emissions.
EIA AEO2006	EIA AEO2006 projects energy supply and demand for the US from 2003 to 2030. Energy consumption is estimated on a regional basis. New Jersey is included in the Middle Atlantic region (NY, PA, CT NJ).	EIA AEO2006 is used to project changes in per capita (residential), per employee (commercial/industrial).
EIA State Electricity Profiles	EIA provides information on the electric power industry generation by primary energy source for 1990 – 2005.	EIA State Electricity Profiles were used to determine the mix of in-state electricity generation by fuel.
US Department of Transportation (DOT), Office of Pipeline Safety (OPS)	Natural gas transmission pipeline mileage, and distribution pipeline mileage and number of services for 1990 – 2005.	Emissions projected to increase at an average annual rate of 0.38% from 2005-2010; 0.45% from 2010-2015; and 0.35% from 2015-2020; based on AEO2006 regional forecast in natural gas consumption for Middle Atlantic region.
NJDEP	NJDEP provided landfill emplacement and control data for New Jersey landfills and CH ₄ emission and wastewater flow data from municipal wastewater facilities.	Waste emplacement data used to estimate emissions from solid waste management. CH ₄ and wastewater data used to estimate wastewater emissions.
US Forest Service; NJDEP; and Center for Remote Sensing and Spatial Analysis, at Rutgers University	Data on forest carbon stocks and land use cover for multiple years.	Data are used to calculate CO ₂ flux over time (terrestrial CO ₂ sequestration in forested areas).
USDS National Agricultural Statistics Service (NASS)	USDA NASS provides data on crops and livestock.	Crop production data used to estimate agricultural residue and agricultural soils emissions; livestock population data used to estimate manure and enteric fermentation emissions.

For electricity, we estimate, in addition to the emissions due to fuels combusted at electricity plants in the State, the emissions related to electricity *consumed* in New Jersey. This entails

accounting for the electricity sources used by New Jersey utilities to meet consumer demands. As this analysis is refined in the future, one could also attempt to estimate other sectoral emissions on a consumption basis, such as accounting for emissions from transportation fuel used in New Jersey, but purchased out-of-state. In some cases, this can require venturing into the relatively complex terrain of life-cycle analysis. In general, we recommend considering a consumption-based approach where it will significantly improve the estimation of the emissions impact of potential mitigation strategies. For example re-use, recycling, and source reduction can lead to emission reductions resulting from lower energy requirements for material production (such as paper, cardboard, and aluminum), even though production of those materials, and emissions associated with materials production, may not occur within the State.

Details on the methods and data sources used to construct the inventories and forecasts for each source sector are provided in the following appendices:

- Appendix A. Electricity Use and Supply
- Appendix B. Residential, Commercial, and Industrial (RCI) Fuel Combustion
- Appendix C. Transportation Energy Use
- Appendix D. Industrial Processes
- Appendix E. Fossil Fuel Extraction and Distribution Industry
- Appendix F. Agriculture
- Appendix G. Waste Management
- Appendix H. Forestry

Appendix A. Energy Supply

Overview

This appendix describes the data sources, key assumptions, and the methodology used to develop an inventory of greenhouse gas (GHG) emissions over the 1990-2004 period associated with the generation of electricity to meet electricity demand in New Jersey. It also describes the data sources, key assumptions, and methodology used to develop a forecast of GHG emissions over the 2005-2020 period associated with meeting future electricity demand in the state under a “business-as-usual (BAU)” scenario.

Emissions (Business-as-Usual)

Total New Jersey Demand for Electricity

Figure A1 shows the historical trend in retail sales of electricity to the RCI sectors in New Jersey for 1990 through 2005. Overall, total retail sales of electricity for all sectors increased by an average annual rate of 1.8% from 1990 to 2005, and by 3.2% from 2000 to 2005. From 1990 through 2005, residential and commercial sales have increased at about the same average annual rate (i.e., 2.6% and 2.5%, respectively) while average annual sales of electricity for the industrial sector have declined slightly (-1.6%). Table A1 shows the values used to develop Figure A1. Table A1 also shows electricity sales from in-state generators of electricity as well as sales of imported electricity to meet New Jersey’s electricity demand.

Figure A1. Electricity Generation to Meet New Jersey Demand (GWh/Year)




Table A1. Electricity Sales to Meet New Jersey Demand (Megawatt-hours per Year (MWh/Year))

Year	Total Demand by New Jersey Consumers ^a	Net In-State Generation ^a	Imported Generation ^b
1990	62,857,000	39,968,834	22,888,166
1991	64,682,551	43,304,711	21,377,840
1992	63,122,000	43,664,014	19,457,986
1993	65,620,646	48,255,318	17,365,328
1994	66,258,000	48,435,342	17,822,658
1995	66,754,000	45,098,658	21,655,342
1996	66,889,430	37,663,185	29,226,245
1997	65,915,000	41,756,261	24,158,739
1998	68,161,512	53,666,002	14,495,510
1999	73,140,489	56,994,767	16,145,722
2000	69,977,000	58,085,215	11,891,785
2001	72,339,691	59,421,254	12,918,437
2002	74,460,421	61,569,387	12,891,034
2003	76,382,512	57,399,351	18,983,161
2004	77,593,167	55,882,342	21,710,825
2005	81,896,813	60,549,582	21,347,231

Note: The generation data reported in this table have not been adjusted to account for electricity transmission and distribution losses (about 7% for New Jersey) and energy used to operate electricity generating facilities.

^a Data from EIA State Electricity Profiles Back Issues - Electricity Information by State,

http://www.eia.doe.gov/cneaf/electricity/st_profiles/backissues.html. For 1990, 1992, 1994, 1995, 1997, and 2000, data from EIA State Electricity Profiles, Table 8 (Retail Sales, Revenue, and Average Retail Price by Sector, 1990 Through 2005) http://www.eia.doe.gov/cneaf/electricity/st_profiles/e_profiles_sum.html. Net in-state generation for 1992, 1994, and 1997 from Table 5 (Electric Power Industry Generation by Primary Energy Source, 1990 Through 2005 (Megawatthours)) of EIA State Electricity Profiles.

^b Imported generation = total generation demand minus net in-state generation.

In-State Generation of Electricity

Emissions associated with the generation of electricity by electricity generating units in New Jersey were estimated using the March 2007 release of the United States Environmental Protection Agency's (US EPA) State Greenhouse Gas Inventory Tool (SGIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for fuel combustion.²¹ The default data used in SGIT for New Jersey are from the United States Department of Energy (US DOE) Energy Information Administration's (EIA) *State Energy Data* (SED) for 1990 through 2003. The SGIT files were updated to include (1) 2004 SED information for New Jersey for all fuel types,²² and (2) 2005 SED information for natural gas and wood

²¹ GHG emissions were calculated using SGIT, with reference to *EIIP, Volume VIII: Chapter 1 "Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels"*, August 2004, and Chapter 2 "Methods for Estimating Methane and Nitrous Oxide Emissions from Stationary Combustion", August 2004. The March 2007 versions of the SGIT Excel files (CO2FFC Module.xls and Stationary Combustion Module.xls) containing SED through 2003 for all fuels and sectors were used as the starting point for preparing the inventory for New Jersey.

²² EIA *State Energy Data 2004*, Complete 2004 data released June 1, 2007.

(http://www.eia.doe.gov/emeu/states/state.html?q_state_a=nj&q_state=NEW%20JERSEY).

consumption.^{23, 24} In addition, SGIT procedures were used, based on New Jersey total municipal solid waste (MSW) combusted,²⁵ to estimate the emissions from New Jersey MSW incinerators.

Figure A2 shows the various types of fuels used by electricity generators into New Jersey for 1990 through 2005. Historically, nuclear energy has been a major source of electricity in New Jersey, followed by natural gas, coal, and petroleum. Electricity generated from nuclear fuel varied significantly from 1990 through 2005, ranging from a low of 29% in 1996 to a high of 59% in 1990. From 2000 through 2005, electricity generated from nuclear fuel remained fairly stable in New Jersey's total fuel mix ranging from 48% to 52% over this six-year period.

Electricity generated from natural gas also varied significantly from 1990 through 2005, ranging from a low of 17% in 1990 to a high of 44% in 1996. From 2000 through 2005, electricity generated from natural gas ranged from 25% to 31% of New Jersey's total fuel mix. Electricity generated from coal from 1990 through 2005 ranged from a low of 11% in 1993 to a high of 21% in 1996 and 1997. From 2000 through 2005, electricity generated from coal ranged from 16% to 19% of New Jersey's total fuel mix. Sales of electricity generated from natural gas and coal peaked from 1995 through 1997 offsetting a decline in sales associated with electricity generated from nuclear fuel during this period. Electricity generated from petroleum from 1990 through 2005 ranged from a low of 1% in 1998, 1999, and 2002 to a high of 5% in 1990 and 1991. From 2000 through 2005, electricity generated from petroleum ranged from 1% to 3% of New Jersey's total fuel mix. Other fuel resources that have been used to generate electricity in New Jersey include industrial gases, landfill gas, MSW, and a small amount of hydroelectric power and pumped storage energy.

Imported Electricity

Emissions associated with the generation of electricity that is imported to New Jersey were estimated using data from the EIA *State Electricity Profiles*.²⁶ The calculation took the total retail electricity sales from the *Profiles* and subtracted from it the net in-state generation, as also reported in the *Profiles* (see Table A1). The difference was then multiplied by the factor of 1,261.7 pounds (lbs) CO₂ per Megawatt-hour (MWh),²⁷ and then augmented with an additional 7% to account for transmission and distribution (T&D) losses.²⁸ The resulting quantity was

²³ EIA *Natural Gas Navigator* (http://tonto.eia.doe.gov/dnav/ng/ng_cons_sum_dcu_SNJ_a.htm).

²⁴ Wood consumption for 2005 (http://www.eia.doe.gov/emeu/states/_seds_updates.html).

²⁵ NJDEP, 2007, personal communication, Michael Aucott, NJDEP, For 1990, total MSW combusted was 430,000 short tons. For 2003, total MSW combusted was 2,149,828 short tons as a result of new MSW combustion capacity brought on-line since 1990. MSW combustion was assumed to be essentially the same for all years from 1995 through 2020.


²⁶ EIA *State Electricity Profiles 2004*, May 2006, and earlier editions, from Energy Information Administration, Office of Coal, Nuclear and Alternate Fuels, U.S. Department of Energy, Washington, DC 20585-0650, http://www.eia.doe.gov/cneaf/electricity/st_profiles/e_profiles_sum.html, accessed September, 2007

²⁷ CEEEP, 2007, Based on emissions rate of the PJM power control area, the primary source of imported electricity in NJ, as reported by Center for Energy, Economic & Environmental Policy (CEEEP), Edward J. Bloustein School of Planning and Public Policy, Rutgers, The State University of New Jersey, 33 Livingston Ave., Room 154, New Brunswick, NJ 08901, June, 2007.

²⁸ As estimated by Bill Dougherty, CCS, personal communication, July 19, 2007, and Julia Hutchins, U.S. DOE EIA, personal communication, July, 2007.

further augmented by multiplying by the factor of 1.006304 to represent additional emissions of CH₄ and N₂O.²⁹

Figure A2. New Jersey In-State Generation by Fuel (1990-2005)



Source: EIA, State Electricity Profiles, Table 5 (Electric Power Industry Generation by Primary Energy Source, 1990 Through 2005 (Megawatt hours)) of EIA State Electricity Profiles, http://www.eia.doe.gov/cneaf/electricity/st_profiles/e_profiles_sum.html.

Table A2. Forecast of New Jersey’s Electricity Demand (MWh/Year)

Year	Scenario ^a	Total Demand by New Jersey Consumers	Net In-State Generation	Imported Generation
2010	BAU	95,152,991	55,313,643	39,839,348
2015	BAU	104,941,739	61,301,353	43,640,386
2015	Potential reductions	87,767,505	58,130,724	29,636,781
2020	BAU	109,054,978	76,060,491	32,994,487
2020	Potential reductions	75,462,324	63,473,994	11,988,330

^a BAU – business as usual.

Reference Case Projections (2005-2020, Business-as-Usual)

Rutgers University’s Center for Energy, Economic & Environmental Policy (CEEPP) provided forecasts of electricity generation to meet New Jersey’s electricity demand for 2010, 2015, and

²⁹ This value was developed by M. Aucott, NJDEP, using the factors of 6×10^{-1} metric tons N₂O per trillion British thermal unit (TBtu) and 10.02 metric tons CH₄ per TBtu for all fuels except natural gas, 0.09×10^{-1} metric tons N₂O per TBtu and 4.75 metric tons CH₄ per TBtu for natural gas, 310 and 21 for the global warming potential of these two gases, respectively, as weighted based on estimated fuel mix of the PJM generation facilities being 15% natural gas with the rest of the fossil fuel being oil and coal.

2020 for a BAU scenario and a scenario representing potential reductions (see Table A2).³⁰ These estimates included those from MSW incinerators, which are connected to the electricity grid in New Jersey. Table A3 shows the annual growth rates from 2005 to 2020 associated with the BAU and potential reductions scenarios.

Table A3. Annual Growth Rates from 2005 to 2020 (%)

Scenario	Year	Total Demand by New Jersey Consumers	Net In-State Generation	Imported Generation
BAU	2005-2010	3.0	-1.8	13.3
BAU	2010-2015	2.0	2.1	1.8
BAU	2015-2020	0.8	4.4	-5.4
Potential reductions	2010-2015	-1.6	1.0	-5.7
Potential reductions	2015-2020	-6.4	0.7	-22.8

The CEEEP has developed a detailed energy modeling system for New Jersey BPU to support development of New Jersey’s Energy Master Plan as well as to support sensitivity analyses of various energy supply and demand scenarios for the state. The modeling approach includes dispatch modeling to identify generation capacity that will be added by in-state generators as well as imported from generators out-of-state to meet New Jersey’s future demand for energy.

Results


Emissions Associated with Business As Usual Scenario

Figure A3 presents the emission inventory and reference case projections for the electricity supply sector. Figure A4 compares emissions on a production (in-state generation) and consumption (in-state generation plus imports) basis. Since New Jersey has been and is projected to continue to be a net importer of electricity, emissions are higher on a consumption basis than on a production basis. Figures A3 and A4 were developed from the emissions data in Table A4. Table A5 shows the relative contributions of emissions associated with each fuel type to total electricity supply sector emissions.

On a consumption basis, emissions associated with the electricity sector are estimated to be about 26.5 MMtCO_{2e} in 1990, with emissions associated with imported electricity accounting for about 53% of total emissions. Emissions associated with the use of coal and natural gas to generate electricity accounted for about 40% of total emissions in 1990, and petroleum and other gases accounted for another 9% of total emissions. In 2000, emissions associated with electricity imports reached a low of about 27% of total of electricity sector emissions, while emissions associate with the use of coal and natural gas accounted for a high of about 66% of electricity sector total emissions.

³⁰ CEEEP, 2007


Figure A3. Electricity Generation GHG Emissions by Fuel Type (1990-2020)



Source: Results in table based on approach described in text.

“Other” includes emissions associated with electricity generated in-state by nuclear, municipal solid waste, and landfill gas, and industrial process gas fuels.

Figure A4. Electricity Generation GHG Emissions – Production and Consumption Basis (1990-2020)



Source: Results in table based on approach described in text.

Table A4. Electricity Generation Emissions Inventory and Reference Case Projections (MMtCO₂e)

Fuel Type	1990	1995	2000	2003	2004	2005	2010	2015	2020
Electricity - Consumption Based	26.5	31.7	27.6	31.0	33.6	35.1	39.1	44.2	43.1
Net Imported Electricity	14.1	13.3	7.3	11.7	13.4	14.8	24.5	26.9	20.3
Electricity - Production Based	12.4	18.3	20.2	19.3	20.3	20.3	14.6	17.3	22.8
Coal	6.93	7.47	10.67	9.94	10.48	9.59	7.42	7.60	8.85
CO ₂	6.89	7.44	10.62	9.89	10.43	9.54	7.38	7.56	8.80
CH ₄ and N ₂ O	0.0335	0.0362	0.0522	0.0486	0.0513	0.0495	0.0383	0.0392	0.0457
Natural Gas	3.64	8.32	7.41	7.15	7.77	8.32	5.49	8.18	12.39
CO ₂	3.63	8.32	7.40	7.14	7.76	8.28	5.46	8.14	12.33
CH ₄ and N ₂ O	0.0034	0.0077	0.0069	0.0067	0.0072	0.0400	0.0262	0.0393	0.0597
Oil	1.70	1.21	0.85	0.93	0.71	1.14	0.42	0.25	0.29
CO ₂	1.70	1.21	0.85	0.93	0.71	1.13	0.42	0.25	0.29
CH ₄ and N ₂ O	0.0055	0.0040	0.0028	0.0030	0.0023	0.0041	0.0015	0.0009	0.0010
Wood (CH ₄ and N ₂ O)	0.01	0.03	0.03	0.02	0.02	0.00	0.00	0.00	0.00
MSW/LFG (CO ₂ , CH ₄ , N ₂ O)	0.17	1.27	1.27	1.27	1.27	1.27	1.27	1.27	1.27
Nuclear (CO ₂ , CH ₄ , N ₂ O)	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.00	0.01

Source: Results in table based on approach described in text.

Electricity consumption emissions are the sum of emissions associated with electricity production and net imported electricity.

MSW - municipal solid waste, LFGTE - Landfill Gas Collection System; these are emissions associated with MWS and LFGTE projects that generate and sell electricity to the power grid.

Table A5. Electricity Sector Proportions of Total Emissions by Fuel Type – Consumption Basis (%)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	26.1	23.6	38.7	27.3	19.0	17.2	20.5
Natural Gas	13.7	26.3	26.9	23.7	14.0	18.5	28.7
Petroleum	6.4	3.8	3.1	3.2	1.1	0.6	0.7
Wood	0.02	0.08	0.10	0.00	0.00	0.00	0.00
Other	0.6	4.0	4.6	3.6	3.2	2.9	3.0
Imported Electricity	53.2	42.2	26.6	42.2	62.7	60.9	47.1

Source: Results in table based on approach described in text.

Note: The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table A4.

During the period from 2005 through 2020, production-based emissions are projected to increase by about 2.5 MMtCO₂e reflecting an annual average increase of about 0.8%. Consumption-based emissions are projected to increase by about 8.0 MMtCO₂e from 2005 through 2020, resulting in an annual average increase of about 1.4%. Thus, imported electricity is projected to be the largest source of emissions increasing by about 5.5 MMtCO₂e from 2005 through 2020. Emissions associated with imported electricity are projected to account for over 60% of New Jersey’s total emissions for this sector from 2010 to 2015, but are projected to decline to about 47% of total emissions in 2020 (see Table A5). From 2005 to 2020, coal and natural gas together are projected to account from 33% to over 50% of total of electricity sector emissions depending on the year.

Appendix B. Residential, Commercial, and Industrial (RCI) Fuel Combustion

Overview

Activities in the RCI³¹ sectors produce carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) emissions when fuels are combusted to provide space heating, water heating, process heating, cooking, and other energy end-uses. Carbon dioxide accounts for over 99% of these emissions on a million metric tons (MMt) of CO₂ equivalent (CO₂e) basis in New Jersey. Direct use of oil, natural gas, coal, and wood in the RCI sectors accounted for an estimated 46 MMtCO₂e of gross greenhouse gas (GHG) emissions in 2005.³²

The following discusses the data sources, methods, assumptions, and results used to construct the inventory (1990 to most recent year for which fuel use data are available (i.e., 2004 or 2005)), and reference case projections (2004 or 2005 to 2020) based on business-as-usual assumptions.

Emissions and Reference Case Projections (Business-as-Usual)

Emissions from direct fuel use were estimated using the March 2007 release of the United States Environmental Protection Agency's (US EPA) State Greenhouse Gas Inventory Tool (SGIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for RCI fuel combustion.³³ The default data used in SGIT for New Jersey are from the United States Department of Energy (US DOE) Energy Information Administration's (EIA) *State Energy Data* (SED) for 1990 through 2003. The SGIT files were updated to include (1) 2004 SED information for New Jersey for all fuel types for each of the RCI sectors,³⁴ and (2) 2005 SED information for natural gas and wood consumption.^{35, 36}

Note that the EIIP methods for the industrial sector exclude from CO₂ emission estimates the amount of carbon that is stored in products produced from fossil fuels for non-energy uses. For example, the methods account for carbon stored in petrochemical feedstocks, and in liquefied petroleum gases (LPG) and natural gas used as feedstocks by chemical manufacturing plants (i.e., not used as fuel), as well as carbon stored in asphalt and road oil produced from petroleum. The carbon storage assumptions for these products are explained in detail in the EIIP guidance

³¹ The industrial sector includes emissions associated with agricultural energy use and fuel used by natural gas transmission and distribution (T&D) industry.

³² Emissions estimates from wood combustion include only N₂O and CH₄. Carbon dioxide emissions from biomass combustion are assumed to be "net zero", consistent with US EPA and Intergovernmental Panel on Climate Change (IPCC) methodologies, and any net loss of carbon stocks due to biomass fuel use should be accounted for in the land use and forestry analysis.

³³ GHG emissions were calculated using SGIT, with reference to *EIIP, Volume VIII*: Chapter 1 "Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels", August 2004, and Chapter 2 "Methods for Estimating Methane and Nitrous Oxide Emissions from Stationary Combustion", August 2004. The March 2007 versions of the SGIT Excel files (CO2FFC Module.xls and Stationary Combustion Module.xls) containing SED through 2003 for all fuels and sectors were used as the starting point for preparing the inventory for New Jersey.

³⁴ EIA *State Energy Data 2004*, Complete 2004 data released June 1, 2007.

(http://www.eia.doe.gov/emeu/states/state.html?q_state_a=nj&q_state=NEW%20JERSEY).

³⁵ EIA *Natural Gas Navigator* (http://tonto.eia.doe.gov/dnav/ng/ng_cons_sum_dcu_SNJ_a.htm).

³⁶ RCI wood consumption for 2005 (http://www.eia.doe.gov/emeu/states/_seds_updates.html).

document.³⁷ The fossil fuel types for which the EIIP methods are applied in the SGIT software to account for carbon storage include the following categories: asphalt and road oil, coking coal, distillate fuel, feedstocks (naphtha with a boiling range of less than 401 degrees Fahrenheit), feedstocks (other oils with boiling ranges greater than 401 degrees Fahrenheit), LPG, lubricants, miscellaneous petroleum products, natural gas, pentanes plus,³⁸ petroleum coke, residual fuel, still gas, and waxes. Data on annual consumption of the fuels in these categories as chemical industry feedstocks were obtained from the EIA SED.

Table B1a shows the growth factors that were applied to the emissions calculated from the latest year for which SED were available to forecast emissions to 2020 for the RCI sectors. The factors selected for the forecast were based on an evaluation of annual growth factors calculated from four different sources of data shown in Table B1b. Table B1b also shows historic annual growth rates for energy use by sector and fuel type developed from the SED for New Jersey.

Table B1a. Annual Growth Rates Applied to Forecast Emissions for the RCI Sectors

Sector	2004/2005-2010 ^a	2010-2015	2015-2020	2004-2020	Reference
Residential					
natural gas	1.3%	1.3%	1.3%	1.3%	BPU ^b
petroleum	-3.2%	-3.2%	-3.2%	-3.2%	BPU
wood	-0.6%	-0.9%	-0.4%	-0.3%	AEO2006 Weighted ^c
coal	0.6%	-0.3%	-0.1%	0.1%	AEO2006 Weighted
Commercial					
natural gas	-0.3%	-0.3%	-0.3%	-0.3%	BPU
petroleum	-6.3%	-6.3%	-6.3%	-6.3%	BPU
wood	0.8%	1.0%	0.7%	0.8%	AEO2006 Weighted
coal	0.6%	1.0%	0.7%	0.8%	AEO2006 Weighted
Industrial					
natural gas	-4.5%	-4.5%	-4.5%	-4.5%	BPU
petroleum	0.3%	0.0%	0.2%	-0.03%	AEO2006 Weighted
wood	1.0%	0.7%	1.0%	0.7%	AEO2006 Weighted
coal	-0.5%	-0.2%	-0.1%	-0.6%	AEO2006 Weighted

^a The latest year for which SED were available to calculate emissions was 2004 for coal and petroleum and 2005 for natural gas and wood.

^b Annual growth rates developed by New Jersey Board of Public Utilities (BPU) based on historical trends in natural gas and distillate fuel oil consumption published by the EIA.

^c Annual growth rates calculated from AEO2006 projections for EIA's Mid Atlantic region, adjusted for New Jersey's projected population for the residential sector, projections for non-manufacturing employment for the commercial sector, and projections for manufacturing employment for the industrial sector.

The New Jersey Board of Public Utilities (BPU) has developed compound annual growth rates for natural gas and distillate fuel to forecast consumption of these two fuels as it has worked on the development of its Energy Master Plan.³⁹ The NJ BPU developed the annual growth rates based on an analysis of historical trends in natural gas and distillate fuel use for New Jersey published by the EIA. The NJ BPU annual growth rates for natural gas and distillate fuel were used to forecast emissions for these two fuels for this analysis. The NJ BPU did not prepare

³⁷ EIIP, Volume VIII: Chapter 1 "Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels", August 2004.

³⁸ A mixture of hydrocarbons, mostly pentanes and heavier fractions, extracted from natural gas.

³⁹ Energy Master Plan, <http://www.state.nj.us/bpu/>.

growth estimates for RCI coal or wood consumption, and were in the process of developing an annual growth factor for industrial distillate fuel consumption during the development of this draft forecast. For RCI coal and wood consumption and industrial petroleum consumption, the annual growth rates applied to forecast emissions were calculated from AEO2006, adjusted for New Jersey’s projected population or employment (see Table B1a).

Table B1b. Historical and Projected Average Annual Growth in Energy Use in New Jersey, by Sector and Fuel, 1990-2020

Sector	Historic from EIA/SED (1990-2004) ^a	NJ BPU (2004-2020) ^b	AEO2006 Weighted by NJ Population or Employment (2004-2020) ^c		
			AEO2006 (2004-2020) ^d	AEO2007 (2004-2020) ^e	
Residential					
natural gas	2.1%	1.3%	0.5%	-0.1%	0.1%
petroleum	-1.9%	-3.2%	-0.8%	-1.4%	-0.2%
wood	-3.9%	NA	-0.3%	-1.0%	-0.2%
coal	-10.5%	NA	0.1%	-0.6%	-1.2%
Commercial					
natural gas	2.7%	-0.3%	1.0%	-0.1%	0.3%
petroleum	-7.3%	-6.3%	0.4%	-0.8%	0.1%
wood	-1.4%	NA	0.8%	-0.4%	0.2%
coal	-5.9%	NA	0.8%	-0.4%	-0.3%
Industrial					
natural gas	-1.1%	-4.5%	0.9%	1.7%	0.5%
petroleum	-4.5%	NA	-0.03%	0.8%	-1.3%
wood	-1.7%	NA	0.7%	1.5%	0.01%
coal	-23.7%	NA	-0.6%	0.3%	-0.8%

^a Compound annual growth rates calculated from EIA SED historical consumption by sector and fuel type for New Jersey. Latest year for which EIA SED information was available for each fuel type is 2004 for all fuel types and 2005 for natural gas and wood. Petroleum includes distillate fuel, kerosene, and liquefied petroleum gases for all sectors plus residual oil for the commercial and industrial sectors.

^b Annual growth rates developed by NJ BPU based on historical trends in natural gas and distillate fuel oil consumption published by the EIA. NA-Not available.

^c Annual growth rates calculated from AEO2006 projections for EIA’s Mid Atlantic region, adjusted for New Jersey’s projected population for the residential sector, projections for non-manufacturing employment for the commercial sector, and projections for manufacturing employment for the industrial sector.

^d Annual growth rates calculated from AEO2006 projections for EIA’s Mid Atlantic region.

^e Annual growth rates calculated from AEO2007 projections for EIA’s Mid Atlantic region.

For RCI coal and wood consumption and industrial petroleum consumption, the “AEO2006 Weighted” annual growth rates were applied to forecast emissions. The “AEO2006 Weighted” annual growth rates were developed from AEO2006 projections for EIA’s Mid Atlantic region, adjusted for New Jersey’s projected population for the residential sector, projections for non-manufacturing employment for the commercial sector, and projections for manufacturing employment for the industrial sector. Annual growth rates were calculated from AEO2006 by first dividing regional residential fuel consumption forecast by the regional population forecast and then weighting the regional growth rates by NJ’s population growth. For the commercial and industrial sectors, a similar approach was used by dividing regional fuel consumption forecast by the regional employment forecast and then weighting the regional growth rates by New Jersey’s

employment growth for these sectors. Regional population and employment forecasts were available for AEO2006, but not available for AEO2007 during the development of this draft inventory and forecast.⁴⁰ For the residential sector, New Jersey's annual population growth is expected to average about 0.65% annually between 2004 and 2020.⁴¹ New Jersey's commercial and industrial employment is projected to increase at compound annual rates averaging about 1.2% and -0.8% between 2006 and 2020, respectively.⁴²

For comparison purposes, Table B1b also shows compound annual growth rates calculated by fuel type from the EIA's *Annual Energy Outlook 2006* (AEO2006) and *Annual Energy Outlook 2007* (AEO2007) regional forecasts for the EIA's Mid Atlantic region, which also includes the states of New York and Pennsylvania.^{43, 44} The differences in the growth rates calculated from the AEO2006 and AEO2007 regional forecasts reflect differences in modeling assumptions that may include fuel and electricity price assumptions and technology changes, as well as to structural changes within each sector (such as shifts in subsectoral shares and in energy use patterns).

Results

Figures B1, B2, and B3 show historical and projected emissions for the RCI sectors in New Jersey from 1990 through 2020. These figures show the emissions associated with the direct consumption of fossil and wood fuels. During the period from 1990 through 2020, the residential sector's share of total RCI emissions from direct fuel use was 34% in 1990, increased to 39% in 2000, decreased slightly to 38% in 2005, and is projected to increase to 44% in 2020. The commercial sector's share of total RCI emissions from direct fuel use was 24% in 1990, increased slightly to 25% in 2000, decreased slightly to 23% in 2005, and is projected to decrease to 21% by 2020. The industrial sector's share of total RCI emissions from direct fuel use was 42% in 1990, decreased to 35% in 2000, increased to 38% in 2005, and is projected to decrease slightly to 35% by 2020.

Natural gas consumption accounts for the highest source of emissions for the residential and commercial sectors, while petroleum consumption accounts for the highest source of emissions for the industrial sector. Emissions associated with natural gas consumption to meet RCI demand accounts for about 74% of the emissions for the residential sector, 83% of the emissions for the commercial sector, and 24% of the emissions for the industrial sector, on average, over the 1990 to 2020 time period. From 1990 to 2020, petroleum is the next highest source of emissions for the residential and commercial sectors, accounting, on average, for about 26% and 17% of total

⁴⁰ AEO2006 employment projections for EIA's Mid Atlantic region obtained through special request from EIA (dated September 27, 2006).

⁴¹ CEEEP, 2007, Center for Energy, Economic & Environmental Policy (CEEEP), Edward J. Bloustein School of Planning and Public Policy, Rutgers, The State University of New Jersey, 33 Livingston Ave., Room 154, New Brunswick, NJ 08901 (<http://www.policy.rutgers.edu/cupr/recon/index.php>). Data provided by Nancy Mantell to CCS on March 30, 2007 (file named "Electricity Prelim Data (3-30-07).xls").

⁴² CEEEP, 2007, Center for Energy, Economic & Environmental Policy (CEEEP), Edward J. Bloustein School of Planning and Public Policy, Rutgers, The State University of New Jersey, 33 Livingston Ave., Room 154, New Brunswick, NJ 08901 (<http://www.policy.rutgers.edu/cupr/recon/index.php>). Data provided by Nancy Mantell to CCS on March 30, 2007 (file named "Electricity Prelim Data (3-30-07).xls").

⁴³ EIA AEO2006 with Projections to 2030, <http://www.eia.doe.gov/oiaf/archive/aeo06/index.html>.

⁴⁴ EIA AEO2007 with Projections to 2030, <http://www.eia.doe.gov/oiaf/aeo/index.html>.

emissions, respectively. For the industrial sector, petroleum is the highest source of emissions accounting, on average, for about 75% of total industrial emissions from 1990 to 2020.


Residential Sector

Figure B1 presents the emission inventory and reference case projections for the residential sector. Figure B1 was developed from the emissions data in Table B2a. Table B2b shows the relative contributions of emissions associated with each fuel type to total residential sector emissions.

For the residential sector, emissions from direct fossil fuel use in 1990 were about 15.6 MMtCO_{2e}, and are estimated to increase to about 19.3 MMtCO_{2e} by 2020. Emissions associated with natural gas consumption accounted for about 60% of total residential emissions in 1990, and are estimated to increase to 86% of total residential emissions by 2020. In 1990, petroleum consumption accounted for about 40% of total residential emissions, and is estimated to account for about 14% of total residential emissions by 2020. Residential-sector emissions associated with the use of coal and wood in 1990 were about 0.12 MMtCO_{2e} combined, and accounted for about 0.8% of total residential emissions. By 2020, emissions associated with the consumption of these two fuels are estimated to fall to 0.06 MMtCO_{2e}, accounting for 0.3% of total residential sector emissions by that year. Coal consumption by the residential sector has been and is expected to continue to be negligible from 1990 through 2020.

For the 15-year period 2005 to 2020, residential-sector GHG emissions associated with the use of natural gas are expected to increase at average annual rate of about 1.6%. Emissions associated with the use of petroleum and wood are expected to decline annually by about -3.6% and -0.6%, respectively. Total GHG emissions for this sector are projected to increase by an average of about 0.6% annually over the 15-year period.

Figure B1. Residential Sector GHG Emissions from Fuel Consumption (1990-2020)



Source: Results in graph based on approach described in text.

Note: Emissions associated with coal and wood combustion are too small to be seen on this graph.

Table B2a. Residential Sector Emissions Inventory and Reference Case Projections (MMtCO₂e)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	0.007	0.002	0.002	0.002	0.002	0.002	0.002
Petroleum	6.18	5.60	4.95	4.53	3.77	3.14	2.62
Natural Gas	9.34	10.69	12.10	13.08	14.16	15.32	16.58
Wood	0.12	0.10	0.06	0.06	0.06	0.06	0.06
Total	15.64	16.40	17.12	17.67	17.99	18.52	19.26

Source: Results in table based on approach described in text.

Table B2b. Residential Sector Proportions of Total Emissions by Fuel Type (%)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Petroleum	39.5%	34.2%	28.9%	25.6%	21.0%	17.0%	13.6%
Natural Gas	59.7%	65.2%	70.7%	74.0%	78.7%	82.7%	86.1%
Wood	0.7%	0.6%	0.4%	0.3%	0.3%	0.3%	0.3%

Source: Results in table based on approach described in text.

Note: The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table B2a.


Commercial Sector

Figure B2 presents the emission inventory and reference case projections for the commercial sector. Figure B2 was developed from the emissions data in Table B3a. Table B3b shows the relative contributions of emissions associated with each fuel type to total commercial sector emissions.

For the commercial sector, emissions from direct fossil fuel use in 1990 were about 11 MMtCO₂e, and are estimated to decrease to about 9.3 MMtCO₂e by 2020. Emissions associated with natural gas consumption accounted for about 57% of total commercial emissions in 1990, and are estimated to increase to 95% of total commercial emissions by 2020. In 1990, petroleum consumption accounted for about 42% of total commercial emissions and is estimated to account for about 5% of total commercial emissions by 2020. Commercial-sector emissions associated with the use of coal and wood in 1990 were about 0.04 MMtCO₂e combined, and accounted for about 0.4% of total commercial emissions. By 2020, emissions associated with the consumption of these three fuels are estimated to be 0.01 MMtCO₂e and to account for 0.1% of total commercial sector emissions.

For the 15-year period 2005 to 2020, commercial-sector GHG emissions associated with the use of natural gas and petroleum are expected to decrease at average annual rates of about -0.3% and -7.7% respectively. Emissions associated with the use of wood and coal use are expected to be negligible. Total GHG emissions for this sector decrease by an average of about -0.95% annually over the 15-year period.

Figure B2. Commercial Sector GHG Emissions from Fuel Consumption (1990-2020)



Source: Results in graph based on approach described in text.

Note: Emissions associated with coal and wood combustion are too small to be seen on this graph.

Table B3a. Commercial Sector Emissions Inventory and Reference Case Projections (MMtCO₂e)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	0.03	0.02	0.01	0.01	0.01	0.01	0.01
Petroleum	4.64	2.42	2.26	1.410	0.94	0.63	0.42
Natural Gas	6.29	7.64	8.73	9.30	9.15	9.00	8.86
Wood	0.01	0.01	0.01	0.00	0.00	0.00	0.00
Total	10.97	10.09	11.01	10.72	10.11	9.65	9.29

Source: Results in table based on approach described in text.

Table B3b. Commercial Sector Proportions of Total Emissions by Fuel Type (%)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	0.2%	0.2%	0.1%	0.1%	0.1%	0.1%	0.1%
Petroleum	42.3%	24.0%	20.6%	13.2%	9.3%	6.5%	4.5%
Natural Gas	57.4%	75.7%	79.2%	86.7%	90.6%	93.3%	95.3%
Wood	0.1%	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%

Source: Results in table based on approach described in text.

Note: The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table B3a.

Industrial Sector


Figure B3 presents the emission inventory and reference case projections for the industrial sector. Figure B3 was developed from the emissions data in Table B4a. Table B4b shows the relative contributions of emissions associated with each fuel type to total industrial sector emissions.

For the industrial sector, emissions from direct fuel use in 1990 were about 19.7 MMtCO₂e and are estimated to decrease to about 15.6 MMtCO₂e by 2020. Emissions associated with natural gas consumption demand accounted for about 24% of total industrial emissions in 1990, and are estimated decline to about 13% of total industrial emissions by 2020. In 1990, petroleum consumption accounted for about 72% of total industrial emissions, and is estimated to account for about 87% of total industrial emissions by 2020. Industrial-sector emissions associated with the use of coal and wood in 1990 were about 0.7 MMtCO₂e combined, and accounted for about 3.4% of total industrial emissions. For 2020, emissions associated with the consumption of these three fuels are estimated to be 0.02 MMtCO₂e, and to account for 0.1% of total industrial sector emissions.

As shown in Figure B3, there was a significant decline in industrial natural gas consumption between 1995 and 2000. To evaluate this trend further, Figure B4 shows GHG emissions associated with industrial fuel consumption for 1990 through 2005 for each fuel type separately and for all fuel types combined. As shown in Figure B4, the significant decline in natural gas consumption actually occurred between 1999 and 2000 and natural gas consumption then continued to decline slightly from 2000 through 2005. The decline in natural gas consumption is partially offset by an increase in petroleum consumption starting in 2000; however, overall consumption of fuels by the industrial sector remained well below 1999 levels. This trend in emissions which is directly related to fuel consumption reflects an overall decline in fuel use by the industrial sector reflecting the economic recession that occurred in the US after 1999. In addition, the decline in natural gas consumption may reflect the industrial sectors response to

high natural gas prices starting in 2000 by increasing its use of petroleum. Note that the EIA was contacted to obtain information to help explain these trends but the EIA was unable to provide any information within the time frame for completing this initial draft inventory.

Figure B3. Industrial Sector GHG Emissions from Fuel Consumption (1990-2020)



Source: Results in graph based on approach described in text.

Note: Emissions associated with wood combustion are too small to be seen on this graph.

Table B4a. Industrial Sector Emissions Inventory and Reference Case Projections (MMtCO₂e)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	0.66	0.03	0.02	0.02	0.02	0.02	0.02
Petroleum	14.17	12.89	10.46	13.62	13.75	13.79	13.66
Natural Gas	4.82	11.24	4.72	3.98	3.14	2.49	1.97
Wood	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Total	19.66	24.16	15.21	17.62	16.91	16.30	15.65

Source: Results in table based on approach described in text.

Table B4b. Industrial Sector Proportions of Total Emissions by Fuel Type (%)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Coal	3.4%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%
Petroleum	72.1%	53.3%	68.8%	77.3%	81.3%	84.6%	87.3%
Natural Gas	24.5%	46.5%	31.0%	22.6%	18.6%	15.3%	12.6%
Wood	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%


Source: Results in table based on approach described in text.

Note: The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table B4a.

For the 15-year period 2005 to 2020, industrial-sector GHG emissions associated with the use of natural gas are expected to decrease at an average annual rate of about -4.6%, and emissions associated with the use of petroleum are expected to increase at an average annual rate of about 0.02%. Emissions associated with the use of wood and coal use are expected to be negligible.

Total GHG emissions for the industrial sector decrease by an average of about -0.8% annually over the 15-year period.

Figure B4. Industrial Sector GHG Emissions from Fuel Consumption (1990-2005)



Source: Results in graph based on approach described in text.

Key Uncertainties

Key sources of uncertainty underlying the estimates above are as follows:

- Population and economic growth are the principal drivers for fuel use. The reference case projections are based on regional fuel consumption projections for EIA's Mid Atlantic modeling region scaled for New Jersey population and employment growth projections. Consequently, there are significant uncertainties associated with the projections. Future work should attempt to base projections of GHG emissions on fuel consumption estimates specific to New Jersey to the extent that such data become available.
- The AEO2006 projections assume no large long-term changes in relative fuel and electricity prices, relative to current price levels and to US DOE projections for fuel prices. Price changes would influence consumption levels and, to the extent that price trends for competing fuels differ, may encourage switching among fuels, and thereby affect emissions estimates.

Appendix C. Transportation Energy Use

Overview

Transportation is the largest greenhouse gas (GHG) source sector in New Jersey –accounting for over one third of New Jersey’s current gross GHG emissions. The transportation sector includes light- and heavy-duty on-road vehicles, aircraft, locomotives, and marine engines. Carbon dioxide (CO₂) accounts for about 97% of the transportation sector’s GHG emissions in 1990 and is projected to increase to about 98% of transportation GHG emissions by 2020. Most of the remaining GHG emissions from the transportation sector are due to nitrous oxide (N₂O) emissions from gasoline engines.

The following discusses the data sources, methods, assumptions, and results used to construct the inventory (1990 to most recent year for which activity data are available (i.e., 2003, 2004, or 2005)), and reference case projections (2004, 2005, or 2006 to 2020) based on business-as-usual assumptions.

Historical Emissions and Reference Case Projections (Business-as-Usual)

Historical greenhouse gas emissions were estimated using the United States Environmental Protection Agency’s (US EPA) State Greenhouse Gas Inventory Tool (SGIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for the sector.^{45,46} In some cases, the SGIT default data or methodologies were refined to account for data or issues specific to New Jersey. Emissions were calculated from 1990 through the most recent year for which the historical activity data were available. This varied from 2003 to 2005, depending upon the data source. For all transportation sectors, the CO₂ emission factors are in units of pounds (lb) per million British thermal unit (MMBtu). The methane (CH₄) and N₂O emission factors are both in units of grams per vehicle mile traveled (VMT) for on-road vehicles and in grams per kilogram of fuel for the aviation, marine vessel, and rail sectors. Key assumptions in this analysis are listed in Table C1. The default fuel consumption data within SGIT were used in most cases to estimate emissions, with the most recently available fuel consumption data (2004) from the United States Department of Energy (US DOE) Energy Information Administration’s (EIA) *State Energy Data* (SED) added.⁴⁷

⁴⁵ CO₂ emissions were calculated using SGIT, with reference to Emission Inventory Improvement Program, Volume VIII: Chapter. 1. “Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels”, August 2004.

⁴⁶ CH₄ and N₂O emissions were calculated using SGIT, with reference to Emission Inventory Improvement Program, Volume VIII: Chapter. 3. “Methods for Estimating Methane and Nitrous Oxide Emissions from Mobile Combustion”, August 2004.

⁴⁷ Energy Information Administration, State Energy Consumption, Price, and Expenditure Estimates (SED), http://www.eia.doe.gov/emeu/states/_seds.html

Table C1. Key Assumptions and Methods for the Transportation Inventory and Projections

Vehicle Type and Pollutants	Methods
<p>On-road gasoline, diesel, and natural gas vehicles and lubricants – CO₂</p>	<p>Inventory (1990 – 2004) EPA SGIT and fuel consumption from EIA SED</p> <p>Reference Case Projections (2005-2020) On-road gasoline and diesel fuel consumption projections, used in the CO₂ emission calculations, are based on VMT projections adjusted by fuel efficiency improvement projections from EIA AEO2006. Other on-road fuels and lubricants projected using Middle Atlantic Region fuel consumption projections from EIA AEO2006, and population projection from SIT Projection tool.</p>
<p>On-road gasoline and diesel vehicles – CH₄ and N₂O</p>	<p>Inventory (1990 – 2005) EPA SGIT emission factors and default EPA SGIT VMT from FHWA by vehicle type.</p> <p>Reference Case Projections (2006-2020) From VMT projections based on NJDOT data from 1990-2005 forecasted to 2020 allocated to vehicle type using EIA AEO2006 VMT projection data.</p>
<p>Aviation – CO₂</p>	<p>Inventory (1990 – 2004) EPA SGIT and fuel consumption from EIA SED. EIA aircraft fuel consumption is adjusted to eliminate the portion of the fuel consumed in New York and the portion used for international flights, based on airport operations from Federal Aviation Administration (FAA) for NJ and NY and fuel consumption for NJ and NY</p> <p>Reference Case Projections (2005-2020) Growth rates are based on projected aircraft operations in New Jersey from FAA, adjusted for fuel efficiency improvements from AEO2006.</p>
<p>Commercial Marine Vessels (residual and diesel) and Recreational Gasoline Vessels – CO₂</p>	<p>Inventory (1990 – 2004) EPA SGIT and fuel consumption from EIA SED for gasoline. National commercial marine fuel consumption from EIA, excluding international bunker fuel, allocated to New Jersey based on New Jersey port tonnage. Offshore emissions pulled from Commission for Environmental Cooperation in North America (CEC) inventory for 2002, allocated to other years based on underway fuel consumption.</p> <p>Reference Case Projections (2005 – 2020) All marine categories projected based on linear regression of historical trend in New Jersey marine fuel consumption.</p>
<p>Non-highway fuel consumption (jet aircraft, aviation gasoline, boats, locomotives) –CH₄, and N₂O</p>	<p>Inventory (1990 – 2003 for boats and aviation; 1990 – 2005 for locomotives) EPA SGIT and fuel consumption from EIA SED.</p> <p>Reference Case Projections (2004 – 2020 for boats and aviation; 2006 – 2020 for locomotives) Aircraft projected using aircraft operations projections from FAA and jet fuel efficiency improvement projections from AEO2006. All marine categories projected based on linear regression of historical trend in New Jersey marine fuel consumption. Locomotives estimated to remain at 2005 levels.</p>

On-road Vehicles

The default SGIT VMT data for 1990 through 2005 in SGIT were compared to annual VMT data obtained from the New Jersey Department of Transportation (NJDOT).⁴⁸ The annual VMT totals included with the default SGIT were all within 1% of the annual VMT data obtained from NJDOT, with the exception of 2003 where the NJDOT data were 2% greater than the default Federal Highway Administration⁴⁹ (FHWA) data included in SGIT. Because the NJDOT VMT totals were almost identical to the SGIT default VMT totals and since the NJDOT data were not broken down according to the vehicle classes needed for the SGIT, the SGIT default VMT data were used in the CH₄ and N₂O emission calculations.

Growth rates for projecting the on-road vehicle gasoline and diesel CH₄ and N₂O emissions were calculated based on a linear regression of the historical 1990 through 2005 VMT data for New Jersey. The total VMT projections were distributed by vehicle type based on national vehicle type VMT forecasts reported in EIA's *Annual Energy Outlook 2006* (AEO2006).⁵⁰ The AEO2006 data were incorporated because they indicate significantly different VMT growth rates for certain vehicle types (e.g., 34% growth between 2002 and 2020 in heavy-duty gasoline vehicle VMT versus 284% growth in light-duty diesel truck VMT over this period). The procedure first applied the AEO2006 vehicle type-based national growth rates to 2005 New Jersey estimates of VMT by vehicle type. These data were then used to calculate the estimated proportion of total VMT by vehicle type in each year. Next, these proportions were applied to the projected estimates for total VMT in New Jersey for each year to yield the vehicle type VMT estimates and compound annual average growth rates are displayed in Tables C2 and C3, respectively. These VMT growth rates were also applied to natural gas vehicles.

Table C2. New Jersey Projected Vehicle Miles Traveled Estimates (millions)

Vehicle Type	2002	2005	2010	2015	2020
Heavy Duty Diesel Vehicle	4,728	4,938	5,539	6,196	6,893
Heavy Duty Gasoline Vehicle	694	729	759	827	901
Light Duty Diesel Truck	691	747	992	1,333	1,803
Light Duty Diesel Vehicle	215	224	298	400	542
Light Duty Gasoline Truck	22,972	24,828	25,959	27,447	28,858
Light Duty Gasoline Vehicle	40,407	42,101	44,017	46,542	48,933
Motorcycle	234	252	264	279	293
Total	69,942	73,819	77,827	83,024	88,222

⁴⁸ New Jersey Department of Transportation, "Roadway Information and Traffic Counts, Public Roadway Mileage and Vehicle Miles Traveled," <http://www.state.nj.us/transportation/refdata/roadway/vmt.shtm>.

⁴⁹ Highway Statistics, Federal Highway Administration, <http://www.fhwa.dot.gov/policy/ohpi/hss/index.htm>.

⁵⁰ "Annual Energy Outlook 2006: With Projections to 2030," U.S. Department of Energy, Energy Information Administration, Washington, DC, DOE/EIA-0383(2006), February 2006. (<http://www.eia.doe.gov/oiaf/aeo06/index.html>)

Table C3. New Jersey Vehicle Miles Traveled Compound Annual Growth Rates

Vehicle Type	2002-2005	2005-2010	2010-2015	2015-2020
Heavy Duty Diesel Vehicle	1.46%	2.32%	2.27%	2.15%
Heavy Duty Gasoline Vehicle	1.65%	0.81%	1.73%	1.72%
Light Duty Diesel Truck	2.62%	5.84%	6.09%	6.23%
Light Duty Diesel Vehicle	1.38%	5.84%	6.09%	6.23%
Light Duty Gasoline Truck	2.62%	0.89%	1.12%	1.01%
Light Duty Gasoline Vehicle	1.38%	0.89%	1.12%	1.01%
Motorcycle	2.51%	0.89%	1.12%	1.01%

For forecasting CO₂ emissions, growth in fuel consumption is needed. On-road gasoline and diesel fuel consumption were forecasted by developing a set of growth factors that adjusted the VMT projections to account for improvements in fuel efficiency. Fuel efficiency projections were taken from AEO2006. The resulting on-road fuel consumption growth rates are shown in Table C4. Growth rates for projecting CO₂ emissions from natural gas vehicles, lubricants, and other fuel consumption were calculated by allocating the AEO2006 consumption of these fuels in the Middle Atlantic region and allocating this to New Jersey based on the ratio of New Jersey's projected population to the Middle Atlantic's projected population.

Table C4. New Jersey On-road Fuel Consumption Compound Annual Growth Rates

Fuel Growth Factors	2002-2005	2005-2010	2010-2015	2015-2020
on-road gasoline	1.84%	1.10%	0.61%	0.95%
on-road diesel	1.33%	2.29%	2.59%	2.50%

Gasoline consumption estimates for 1990 through 2004 were adjusted by subtracting ethanol consumption, per the methodology used in SGIT. The historical EIA ethanol consumption data show that use of ethanol in New Jersey began in 1993, peaked in 2001, and dropped off considerably in 2002 and 2003, and then increased again in 2004 with ethanol consumption ranging from 0.02% to 0.2% of the gasoline consumption on a Btu basis. Within the last several years ethanol consumption has risen again and is currently estimated to represent nearly 10% by volume of New Jersey gasoline sales. For the reference case projections, ethanol consumption was assumed to remain at the 2004 level. Even though current ethanol consumption is higher than this, overall estimates of CO₂ emissions from motor vehicle fuel consumption are not estimated to be affected significantly, since the ethanol presently in the market is derived from corn and differs little from gasoline in overall life cycle carbon emissions.⁵¹

Rail and Recreational Marine Vehicles

For the rail and recreational marine sectors, 1990 through 2004 estimates are based on SGIT methods and fuel consumption from EIA. Marine gasoline consumption was projected to 2020 based on a linear regression of the 1990 through 2005 historical data. The historic data for rail shows no significant positive or negative trend; therefore, no growth was assumed for this sector.

⁵¹ Farrell, et al., 2006, Ethanol can contribute to energy and environmental goals, *Science*, 311, 506-508.

Commercial Marine Vessels

For the commercial marine sector (marine diesel and residual fuel), 1990 through 2004 emission estimates are based on SGIT emission rates applied to estimates of New Jersey marine vessel diesel and residual fuel consumption. Because the SGIT default relies on marine vessel fuel consumption estimates that represent the State in which fuel is sold rather than consumed, an alternative method was used to estimate New Jersey marine vessel fuel consumption. New Jersey fuel consumption estimates were developed by allocating 1990 through 2004 national diesel and residual oil vessel bunkering fuel consumption estimates obtained from EIA.⁵² The annual estimates of the U.S. amount of international marine vessel bunker fuels was subtracted from the 1990 through 2004 national diesel and residual oil vessel bunkering fuel consumption estimates. Marine vessel fuel consumption was allocated to New Jersey using the marine vessel activity allocation methods/data compiled to support the development of EPA’s National Emissions Inventory (NEI).⁵³ In keeping with the NEI, 75% of each year’s distillate fuel and 25% of each year’s residual fuel were assumed to be consumed within the port area, and the remaining fuel consumption was assumed to occur while ships are underway. National port area fuel consumption was allocated to New Jersey based on year-specific freight tonnage data for New Jersey as reported in “Waterborne Commerce of the United States, Part 5 – Waterways and Harbors National Summaries.”⁵⁴ Offshore CO₂ and hydrocarbon (HC) emissions for New Jersey’s exclusive economic zone (EEZ) was taken from a study by Corbett for the Commission for Environmental Cooperation (CEC) in North America.⁵⁵ Offshore CH₄ emissions were estimated by speciating the HC emissions using the California Air Resources Board’s total organic gas (TOG) profile (#818).⁵⁶ Offshore N₂O emissions were estimated by applying the ratio of N₂O to CH₄ emission factors to the CH₄ emission estimate. The 2002 offshore emissions from the CEC inventory were scaled to other historic years based on the estimated underway diesel and residual fuel consumption. Port and offshore commercial marine emissions were projected using linear regression based on the 1990 through 2004 emission data.

The resulting compound annual average growth rates for marine fuel consumption are displayed in Table C5.

Table C5. New Jersey Compound Annual Growth Rates for Marine Fuels

Fuel	2002-2005	2005-2010	2010-2015	2015-2020
Marine Gasoline	8.27%	-1.37%	0.99%	0.94%
Marine Diesel	-1.12%	1.86%	1.70%	1.57%
Marine Residual	-7.25%	1.34%	1.26%	1.18%

⁵² US Department of Energy, Energy Information Administration, “Petroleum Navigator” (diesel data obtained from <http://tonto.eia.doe.gov/dnav/pet/hist/kd0vabnus1a.htm>; residual data obtained from <http://tonto.eia.doe.gov/dnav/pet/hist/kprvatnus1a.htm>).

⁵³ See methods described in

ftp://ftp.epa.gov/EmisInventory/2002finalnei/documentation/mobile/2002nei_mobile_non-road_methods.pdf

⁵⁴ Note that it was necessary to estimate the 1999 and 2000 values by interpolation based on the 1998 and 2001 values and the 1990-1993 values by extrapolating based on the 1994 through 2004 estimates.

⁵⁵ Estimate, Validation, and Forecasts of Regional Commercial Marine Vessel Inventories, submitted by J. Corbett, prepared for the California Air Resources Board, California Environmental Protection Agency, and Commission for Environmental Cooperation in North America, <http://coast.cms.udel.edu/NorthAmericanSTEEM/>.

⁵⁶ California Air Resources Board, Speciation Profiles, <http://www.arb.ca.gov/ei/speciate/speciate.htm>.

Table C6 summarizes the GHG emission totals for the marine sector (including gasoline-, distillate-, and residual-fueled vessels). In this table, the offshore emissions are listed separately. Although this number includes emissions from vessels entering and leaving New Jersey ports, it also includes emissions from ships passing through the area, within 200 miles of the New Jersey shoreline. Because New Jersey has no control over emissions from the vessels passing through the area, these emissions are not included in the emission summary totals.

Table C6. Summary of New Jersey GHG Marine Emissions

Emission Totals (MMtCO₂e)	1990	1995	2000	2005	2010	2015	2020
Boats and Ships - Ports/Inshore	1.0	1.2	1.4	1.5	1.6	1.7	1.8
Boats and Ships - Offshore	1.1	1.9	2.2	1.7	1.8	1.9	2.0

Aviation

For the aircraft sector, emission estimates for 1990 to 2004 are based on SGIT emission factors. The New Jersey fuel consumption data from EIA includes a significant portion of jet fuel and aviation gasoline purchased in New Jersey, but used at New York airports. In addition, the EIA fuel consumption includes a portion of jet fuel consumed for international transport. Thus, the EIA fuel consumption data were adjusted to account for the portion of jet fuel and aviation gasoline actually consumed in New Jersey, and not on international flights. This was done by first calculating an adjustment factor that would account for the portion of fuel actually consumed in New Jersey.

Data on jet fuel and aviation gasoline operations (i.e., landings and takeoffs) by year for New Jersey and New York were obtained from the Federal Aviation Administration’s Terminal Area Forecast System.⁵⁷ The fraction of New Jersey jet fuel operation was calculated as the number of total jet fuel operations in New Jersey divided by the total number of jet fuel operations in New York and New Jersey combined, for each year from 1990 through 2004. Similarly, the fraction of jet fuel consumed in New Jersey was calculated as the EIA jet fuel consumed in New Jersey divided by the total jet fuel consumed in New York and New Jersey for each year. A New Jersey adjustment factor was then calculated for each year from 1990 through 2004 by dividing the New Jersey fraction of jet fuel operations by the fraction of New Jersey jet fuel consumption. This yearly New Jersey adjustment factor was then multiplied by the EIA jet fuel consumption for New Jersey for each year from 1990 through 2004. Similar adjustment factors were also calculated for aviation gasoline using aviation gasoline operations and aviation gasoline consumption.

Next, yearly adjustment factors to exclude the portion of fuel from international flights were calculated. The yearly amount of jet fuel consumed for international transport from the U.S. was calculated by EPA.⁵⁸ This amount was subtracted from the EIA total U.S. jet fuel consumption for each year from 1990 through 2004 and then divided the result by the total yearly U.S. jet fuel

⁵⁷ Terminal Area Forecast, Federal Aviation Administration, <http://www.apo.data.faa.gov/main/taf.asp>.

⁵⁸ “Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2005,” USEPA #430-R-07-002, April 2007, Table 3-50, “US Aviation Jet Fuel Consumption for International Transport,” (<http://www.epa.gov/climatechange/emissions/downloads06/07Energy.pdf>)

consumption. This fraction was then multiplied by the New Jersey jet fuel consumption after the New York/New Jersey adjustment had been applied.

Aviation emissions were projected from 2005 through 2020 using general aviation and commercial aircraft operations projections data from the Federal Aviation Administration’s Terminal Area Forecast System⁵⁹ and national aircraft fuel efficiency forecasts. To estimate changes in jet fuel consumption, itinerant aircraft operations from air carrier, air taxi/commuter, and military aircraft were first summed for each year of interest. The post-2002 estimates were adjusted to reflect the projected increase in national aircraft fuel efficiency (indicated by increased number of seat miles per gallon), as reported in AEO2006. Because AEO2006 does not estimate fuel efficiency changes for general aviation aircraft, forecast changes in aviation gasoline consumption were based solely on the projected number of itinerant general aviation aircraft operations in New Jersey, which was obtained from the Federal Aviation Administration (FAA) source noted above. The resulting compound annual average growth rates are displayed in Table C7. The negative growth for aviation gasoline for the 2002-2005 period is supported by prime supplier sales volumes from EIA, which shows sales of 14.8 thousand gallons per day in 2002 and 12.6 thousand gallons per day in 2005.⁶⁰

Table C7. New Jersey Aviation Fuels Compound Annual Growth Rates

Fuel	2002-2005	2005-2010	2010-2015	2015-2020
Aviation Gasoline	-1.87%	0.34%	1.40%	0.93%
Jet Fuel	0.37%	0.94%	0.53%	0.42%

Because much of the aviation fuel allocated to New Jersey is actually consumed outside of New Jersey’s air space, and cannot be controlled by actions established by New Jersey, NJDEP felt it was more appropriate to account for only the portion of aviation emissions under New Jersey’s control. Based on the above calculations, and rough estimations of the portion of in-state flights, which New Jersey regulations could impact, NJDEP believes that approximately 1.0 MMtCO₂e can be accounted for and controlled by New Jersey actions. Table C8 summarizes the results of the original fuel-based CO₂ calculations, along with New Jersey’s revised assumptions based on the portion of New Jersey-controlled fuel.

Table C8. Summary of New Jersey Aviation GHG Emissions

Emission Totals (MMtCO ₂ e)	1990	1995	2000	2005	2010	2015	2020
Aviation—all NJ-based fuel	4.8	5.4	4.2	4.0	4.2	4.3	4.4
Aviation—NJ controlled fuel	1.0	1.0	1.0	1.0	1.0	1.0	1.0

Non-road Engines

It should be noted that fuel consumption data from EIA includes non-road gasoline and diesel fuel consumption in the commercial and industrial sectors. Emissions from these non-road engines are included in the inventory and forecast for the residential, commercial, and industrial (RCI) sectors. Table C9 shows how EIA divides gasoline and diesel fuel consumption between the transportation, commercial, and industrial sectors.

⁵⁹ Terminal Area Forecast, Federal Aviation Administration, <http://www.apo.data.faa.gov/main/taf.asp>.

⁶⁰ US Department of Energy, Energy Information Administration, “Petroleum Navigator”, <http://tonto.eia.doe.gov/dnav/pet/hist/c400013451a.htm>.

Table C9. EIA Classification of Gasoline and Diesel Consumption


Sector	Gasoline Consumption	Diesel Consumption
Transportation	Highway vehicles, marine	Vessel bunkering, military use, railroad, highway vehicles
Commercial	Public non-highway, miscellaneous use	Commercial use for space heating, water heating, and cooking
Industrial	Agricultural use, construction, industrial and commercial use	Industrial use, agricultural use, oil company use, off-highway vehicles

Results

Figure C1 illustrates the trend in the transportation sector GHG emissions from 1990 through 2020. The results are also listed in Table C10. As shown in Figure C1 and Table C10, on-road gasoline consumption accounts for the largest share of transportation GHG emissions. Emissions from on-road gasoline vehicles increased by about 29% from 1990 to 2004 and contributed 78% of total transportation emissions in 2004. GHG emissions from on-road diesel fuel consumption increased by 79% from 1990 to 2004, and by 2004 accounted for 15% of GHG emissions from the transportation sector. In contrast, emissions from aviation stayed flat between 1990 and 2004 to cover just 2% of transportation emissions in 2002, and port and inshore emissions from boats and ships increased by 47% from 1990 to 2004 to cover 3% of transportation emissions in 2004. Emissions from all other categories combined (locomotives, natural gas, and oxidation of lubricants) contributed just over 1% of total transportation emissions in 2004.

GHG emissions from all on-road vehicles combined are projected to increase by 21% between 2004 and 2020. Historical growth for diesel fuel was stronger than for gasoline. This trend is expected to continue for the 2004-2020 period, with on-road gasoline and diesel fuel emissions projected to increase by 16% and 46%, respectively. Jet fuel and aviation gasoline consumption is projected to continue to remain flat between 2004 and 2020. Growth in marine vessel emissions is projected to slow in the projection years, with an increase of 20% from 2004 to 2020.

Figure C1. New Jersey Transportation GHG Emissions by Sector, 1990-2020



Source: Results in graph based on approach described in text.

Table C10. Summary of New Jersey GHG Emissions and Reference Case Projections

Emission Totals (MMtCO ₂ e)	1990	1995	2000	2005	2010	2015	2020
On-road Gasoline	29.8	31.1	35.6	38.9	41.1	42.4	44.3
On-road Diesel	4.2	4.7	6.8	7.6	8.5	9.7	11.0
Aviation	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Boats and Ships - Ports/Inshore	1.0	1.2	1.4	1.5	1.6	1.7	1.8
Rail	0.3	0.7	0.2	0.2	0.2	0.2	0.2
Other	0.3	0.3	0.3	0.3	0.3	0.4	0.4
Total	36.6	39.1	45.2	49.5	52.7	55.3	58.6

Source: Results in table based on approach described in text.

Reductions from Potential Actions

California Light-Duty GHG Standards

On-road gasoline and diesel emissions in New Jersey were also adjusted to reflect the effects of California's light-duty vehicle GHG standards. Note that the implementation pathway of these standards is not totally certain at this time due to EPA's denial of California's request and subsequent legal action against EPA on the part of several states, including New Jersey. For the purpose of this analysis, these standards were applied to new vehicles starting with model year 2009. First, the projected fuel consumption for new vehicles without the California GHG standards was estimated for light-duty vehicles by applying projected new vehicle fuel economy from AEO2006 to the estimated VMT for each model year from 2009 through 2020. VMT by model year for model year 2009 and newer vehicles was estimated using New Jersey's registration distribution data developed by NJDEP based on a June 4, 2003 snapshot of the New Jersey vehicle registration database, combined with MOBILE6 default annual mileage accumulation rates by model year and vehicle type. CO₂ emissions for these vehicles without standards were then estimated by applying SGIT default CO₂ emission factors to the estimated fuel consumption. CO₂ emissions for the vehicles under the standards were estimated by

applying the emission levels set by the California standards (in CO₂e-g/mi) to the estimated VMT by model year. The emission reductions resulting from the standards were then estimated by subtracting estimated emissions from phased-in light-duty vehicles from the estimated emissions for these vehicles without the standards. By 2020, the California standards resulted in a 22% reduction in CO₂ emissions from on-road gasoline and a 2% reduction in CO₂ emissions from on-road diesel fuel.

Table C11 summarizes emissions from the transportation sector after accounting for potential reduction actions in 2020.

Table C11. Summary of New Jersey GHG Emissions and Reference Case Projections

Emission Totals (MMtCO₂e)	2005	2020 Reference Case Projections	2020 with Potential Reductions
On-road Gasoline	38.9	44.3	31.3
On-road Diesel	7.6	11.0	10.5
Aviation	1.0	1.0	1.0
Boats and Ships - Ports/Inshore	1.5	1.8	1.8
Rail	0.2	0.2	0.2
Other	0.3	0.4	0.4
Total	49.5	58.6	45.2

Key Uncertainties

Uncertainties in On-road Vehicle Fuel Consumption

A major uncertainty in this analysis is the projected increase in on-road VMT and gasoline consumption from 2006 to 2020. The VMT projections are based on linear regressions of the historical 1990 through 2005 VMT data. It would be preferable to obtain VMT growth rates based on in-State VMT projections. Another uncertainty is the future year vehicle mix, which was calculated based on national growth rates for specific vehicle types. These growth rates may not reflect vehicle-specific VMT growth rates for the state.

The growth rate in VMT and fuel consumption from heavy-duty diesel trucks may be higher than that projected here. A study of projected heavy-duty truck traffic in New Jersey suggests that heavy-duty diesel VMT will increase by 80% from 1998 to 2020.⁶¹ This compares to a growth rate in on-road diesel fuel consumption of 68% from 1998 to 2020, used for the GHG emission projections in this analysis. Based on a growth rate of 80%, on-road diesel GHG emissions would be about 11.7 MMtCO₂e in 2020, an increase of 0.8 MMtCO₂e from the results presented here.

Uncertainties in Aviation Fuel Consumption

The jet fuel and aviation gasoline fuel consumption from EIA is actually fuel *purchased* in the State, and therefore, includes fuel consumed during state-to-state flights and international flights. The methodology used here attempts to account for the portion of the fuel consumed in New York and the portion consumed in international flights. However, the methodology used to estimate the portion of fuel consumed on international flights assumes that each State's share of

⁶¹ "The Trucks Are Coming: What Growing Truck Traffic Will Mean for New Jersey's Quality of Life," Tri-State Transportation Campaign, January 2005.

international aviation fuel consumption is the same as the State's share of total aviation jet fuel consumption. It is likely that New Jersey's share of fuel consumption for international flights would be higher than this, indicating that aviation emissions for New Jersey may be overestimated.

Uncertainties in Marine Fuel Consumption

There are several assumptions that introduce uncertainty into the estimates of commercial marine fuel consumption. These assumptions include:

- 75% of marine diesel and 25% of residual fuel is consumed in port;
- The proportion of freight tonnage for New Jersey as a fraction of national freight tonnage appropriately represents marine fuel that is consumed in New Jersey;
- The 2002 estimate of underway emissions for New Jersey includes marine vessel travel 200 miles out to sea. This may actually include marine vessels traveling to New York ports, as well as New Jersey ports, which may lead to an overestimation of the underway marine vessel emissions.

Appendix D. Industrial Processes

Overview

Emissions in the industrial processes category span a wide range of activities, and reflect non-combustion sources of greenhouse gas (GHG) emissions from several industries. The industrial processes that exist in New Jersey, and for which emissions are estimated in this inventory, include the following:

- Carbon Dioxide (CO₂) from consumption of limestone and soda ash;
- Nitrous oxide (N₂O) from nitric acid production;
- Hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) from consumption of substitutes for ozone-depleting substances (ODS) used in cooling and refrigeration equipment;
- HFCs, PFCs, and sulfur hexafluoride (SF₆) from semiconductor manufacture; and
- SF₆ from transformers used in electric power transmission and distribution (T&D) systems and the United States (US) Department of Energy's (DOE) Princeton Plasma Physics Laboratory (PPPL).

Other industrial processes that are sources of GHG emissions but are not found in New Jersey include the following:

- CO₂ from cement manufacture, lime and soda ash production, and dolomite consumption;
- CO₂ and methane (CH₄) from iron and steel production;
- CO₂, CH₄, N₂O from ammonia manufacture and urea application;
- N₂O from adipic acid production;
- PFCs from aluminum production;
- SF₆ from magnesium production and processing; and
- HFCs from HCFC-22 production.

The following discusses the data sources, methods, assumptions, and results used to construct the inventory and reference case projections for this sector. The reference case projections assume business-as-usual practices.

Emissions and Reference Case Projections (Business-as-Usual)

Greenhouse gas emissions for 1990 through 2005 were estimated using the United States Environmental Protection Agency's (US EPA) State Greenhouse Gas Inventory Tool (SGIT) software, and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for this sector.⁶² Table D1 identifies for each emissions source category the information needed for input into SGIT to calculate emissions, the data sources used for the analysis described here, and the historical years for which emissions were calculated based on the availability of data. The New Jersey Department of Environmental Protection (NJDEP) provided data for nitric acid production for 1994 through 2003. NJDEP also obtained data from the US DOE PPPL, formerly a user of relatively significant quantities of SF₆, on SF₆ usage and

⁶² GHG emissions were calculated using SGIT, with reference to EIIP, Volume VIII: Chapter. 6. "Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes", August 2004. Referred to as "EIIP" below.

emissions for 1997 through 2005. Table D1 provides details on how the data provided were used to calculate historical emissions for these three categories.

Table D1. Approach to Estimating Historical Emissions

Source Category	Time Period	Required Data for SGIT	Data Source
Limestone Consumption	1994 - 2003	Mt of limestone consumed.	For default limestone data, the state's total limestone consumption (as reported by the United States Geological Survey (USGS)) is multiplied by the ratio of national limestone consumption for industrial uses to total national limestone consumption. Additional information on these calculations, including a definition of industrial uses, is available in Chapter 6 of the EIIP guidance. Default limestone production data are not available in SGIT for 1990 - 1993. SGIT does not contain default consumption data for dolomite for any year for New Jersey.
Nitric Acid Production	1994 - 2003	Mt of nitric acid produced.	NJDEP provided nitric acid production data from its Release and Pollution Prevention Reports for EI DuPont Denemours & Co., Inc for 1994-2003, and for Saint Gobain Performance Plastics Corp. that produced nitric acid in 1998 only. NJDEP also provided 1991 production data for one plant that was used for calculating 1990 emissions.
Soda Ash Consumption	1990 - 2005	Mt of soda ash consumed for use in consumer products such as glass, soap and detergents, paper, textiles, and food. Emissions based on state's population and estimates of emissions per capita from the US EPA national GHG inventory.	<i>USGS Minerals Yearbook, 2004: Volume I, Metals and Minerals</i> , (http://minerals.usgs.gov/minerals/pubs/commodity/soda_ash/).
ODS Substitutes	1990 - 2005	Based on state's population and estimates of emissions per capita from the US EPA national GHG inventory.	National emissions from US EPA 2007 <i>Inventory of US Greenhouse Gas Emissions and Sinks: 1990-2005</i> (http://www.epa.gov/climatechange/emissions/usinventoryreport.html). US and New Jersey population for 1990-2005 from US Census Bureau (http://www.census.gov).

Table D1. Approach to Estimating Historical Emissions (Continued)

Source Category	Time Period	Required Data for SGIT	Data Source
Semiconductor Manufacturing	1990 - 2005	State and national value of semiconductor shipments for NAICS code 334413 (Semiconductor and Related Device Manufacturing). Method uses ratio of state-to-national value of semiconductor shipments to estimate state's proportion of national emissions for 1990 - 2005.	National emissions from US EPA 2007 <i>Inventory of US Greenhouse Gas Emissions and Sinks: 1990-2005</i> (http://www.epa.gov/climatechange/emissions/usinventoryreport.html). Value of shipments from U.S Census Bureau's 1997 and 2002 <i>Economic Census</i> (http://www.census.gov/econ/census02/). For 1997, value of shipments for state was 0.6% of national total. For 2002, value of shipments for
Electric Power T&D Systems	1990 - 2005	Emissions from 1990 to 2005 based on the national emissions per kWh and state's electricity use provided in SGIT.	National emissions per kWh from US EPA 2007 "Inventory of US Greenhouse Gas Emissions and Sinks: 1990-2005" (http://www.epa.gov/climatechange/emissions/usinventoryreport.html).
Laboratory Use of SF ₆	1990 - 2005	Mt of SF ₆ gas released to atmosphere.	PPPL provided estimates of SF ₆ annual leak rates for 1997-2005.

Table D2 lists the data sources used to quantify activities related to industrial process emissions, the annual compound growth rates implied by estimates of future activity used, and the years for which the reference case projections were calculated.

Table D2. Approach to Estimating Projections

	Time Period	Projection Assumptions	Data Source	Annual Growth Rates (%)			
				2000 to 2005	2005 to 2010	2010 to 2015	2015 to 2020
Limestone Consumption	2004 - 2020	Compound annual employment growth rate for New Jersey's manufacturing sector for 2000-2010. Assumed no change in activity from 2004-2005.	Based on R/ECON modeling forecast of manufacturing employment for 2006-2020. ^a	0.0%	-1.3%	-1.0%	-0.4%
Nitric Acid Production	2004 - 2020	Ditto	Ditto	0.0%	-1.3%	-1.0%	-0.4%
Soda Ash Consumption	2006 - 2020	Growth between 2004 and 2009 is projected to be about 0.5% per year for US production. Assumed growth is same for 2010 – 2020.	<i>Minerals Yearbook, 2005: Volume I, Soda Ash</i> , (http://minerals.usgs.gov/minerals/pubs/commodity/soda_ash/soda_myb05.pdf).	0.5	0.5	0.5	0.5

^a CEEEP, 2007, Center for Energy, Economic & Environmental Policy (CEEEP), Edward J. Bloustein School of Planning and Public Policy, Rutgers, The State University of New Jersey, 33 Livingston Ave., Room 154, New Brunswick, NJ 08901 (<http://www.policy.rutgers.edu/cupr/recon/index.php>). Data provided by Nancy Mantell to CCS on March 30, 2007 (file named "Electricity Prelim Data (3-30-07).xls").

Table D2. Approach to Estimating Projections (Continued)

	Time Period	Projection Assumptions	Data Source	Annual Growth Rates (%)			
				2000 to 2005	2005 to 2010	2010 to 2015	2015 to 2020
ODS Substitutes	2006 - 2020	Based on national growth rate for use of ODS substitutes.	EPA, 2004 ODS substitutes cost study report (http://www.epa.gov/ozone/snap/emissions/TMP6si9htnvca.htm).	15.8	7.9	5.8	5.3
Electric Power T&D Systems	2006 - 2020	Based on national growth rate for use of ODS substitutes.	EPA, 2004 ODS substitutes cost study report (http://www.epa.gov/ozone/snap/emissions/TMP6si9htnvca.htm).	3.3	-6.2	-9.0	-2.8
Laboratory Use of SF ₆	2006-2010	Assumed no growth because SF ₆ gas use and leaks at PPPL facility in the future are unknown.		0.0%	0.0%	0.0%	0.0%

Results


Figures D1 and D2 show historic and projected emissions for the industrial processes sector from 1990 to 2020. Table D3 shows the historic and projected emission values upon which Figures D1 and D2 are based. Total gross New Jersey GHG emissions were about 1.26 MMtCO₂e in 1990, 4.02 MMtCO₂e in 2005, and are projected to increase to about 8.61 MMtCO₂e in 2020. Emissions from the overall industrial processes category are expected to grow by about 5.2% annually from 2005 through 2020, as shown in Figures D1 and D2, with emissions growth primarily associated with increasing use of HFCs and PFCs in refrigeration and air conditioning equipment.

Substitutes for Ozone-Depleting Substances (ODS)

HFCs and PFCs are used as substitutes for ODS, most notably CFCs (CFCs are also potent warming gases, with global warming potentials on the order of thousands of times that of CO₂ per unit of emissions) in compliance with the *Montreal Protocol* and the *Clean Air Act Amendments of 1990*.⁶³ Even low amounts of HFC and PFC emissions, for example, from leaks and other releases associated with normal use of the products, can lead to high GHG emissions on a CO₂e basis. Emissions from the use of ODS substitutes in New Jersey were calculated using the default methods in SGIT (see dark green line in Figure D2). Emissions have increased from 0.01 MMtCO₂e in 1990 to about 2.41 MMtCO₂e in 2000, and are expected to increase to about 8.4 MMtCO₂e in 2020, reflecting an average annual increase of about 6.4% from 2000 to 2020 due to increased substitutions of these gases for ODS. The projected rate of increase for these emissions is based on projections for national emissions from the US EPA report referenced in Table D2.

⁶³ As noted in EIIP Chapter 6, ODS substitutes are primarily associated with refrigeration and air conditioning, but also many other uses including as fire control agents, cleaning solvents, aerosols, foam blowing agents, and in sterilization applications. The applications, stocks, and emissions of ODS substitutes depend on technology characteristics in a range of equipment types. For the US national inventory, a detailed stock vintaging model was used to track ODS substitutes uses and emissions, but this modeling approach is not applied at the state level.

Figure D1. Non-combustion GHG Emissions from Industrial Processes, 1990-2020



Source: Results in graph based on approach described in text.

Table D3. Historic and Projected Emissions for the Industrial Processes Sector (MMtCO₂e)


Industry	1990	1995	2000	2005	2010	2015	2020
Limestone & Dolomite Use (CO ₂)	-	0.013	0.003	0.005	0.005	0.004	0.004
Nitric Acid Production (N ₂ O)	0.203	0.0003	0.001	0.001	0.001	0.001	0.001
Soda Ash Use (CO ₂)	0.084	0.082	0.079	0.076	0.077	0.079	0.081
ODS Substitutes (HFC, PFC)	0.010	0.977	2.413	3.594	5.160	6.818	8.374
Semiconductor Manufacture (PFC; HFC CF ₄ , C ₂ F ₆ , C ₃ F ₈ ; HFC-23; and SF ₆)	0.011	0.019	0.032	0.025	0.018	0.012	0.011
Electricity T&D Systems (SF ₆)	0.629	0.483	0.311	0.296	0.209	0.139	0.124
Laboratory Use of SF ₆	0.325	0.325	0.018	0.018	0.018	0.018	0.018
Total	1.263	1.900	2.857	4.015	5.488	7.072	8.613

Electricity Distribution

Emissions of SF₆ from electrical equipment have experienced declines since the early nineties (see brown line in Figure D2), mostly due to voluntary action by industry. SF₆ is used as an electrical insulator and interrupter in the electricity T&D system. The largest use for SF₆ is as an electrical insulator in electricity T&D equipment, such as gas-insulated high-voltage circuit breakers, substations, transformers, and transmission lines, because of its high dielectric strength and arc-quenching abilities. Not all of the electric utilities in the US use SF₆; use of

the gas is more common in urban areas where the space occupied by electrical distribution and transmission facilities is more valuable.⁶⁴

Figure D2. Non-combustion GHG Emissions from Industrial Processes, 1990-2020, by Source



Source: Results in graph based on approach described in text.

Emissions associated with electricity distribution in New Jersey from 1990 to 2002 were estimated based on the estimates of emissions per kWh of electricity consumed from the US EPA GHG inventory, and the ratio of New Jersey's to the US electricity consumption estimates available from the Energy Information Administration (EIA) and provided in SGIT. The *US Climate Action Report* shows expected decreases in these emissions at the national level, and the same rate of decline is assumed for emissions in New Jersey. The decline in SF₆ emissions in the future reflects expectations of future actions by the electric industry to reduce these emissions. Relative to total industrial non-combustion process emissions, SF₆ emissions from electrical equipment are about 0.63 MMtCO_{2e} in 1990 and 0.12 MMtCO_{2e} in 2020.

Laboratory Use of SF₆

The PPPL used SF₆ in a large fusion experiment that was shutdown in 1997. Information on the amount of SF₆ gas leaks from the large fusion experiment for 1990 through 1997 is not available, but was earlier estimated as being in the range of 28,000 to 36,000 pounds per

⁶⁴ US EPA, Draft User's Guide for Estimating Carbon Dioxide, Nitrous Oxide, HFC, PFC, and SF₆ Emissions from Industrial Processes Using the State Inventory Tool, prepared by ICF International, March 2007.

year.⁶⁵ Since 1997, PPPL has been operating an experiment that requires much less SF₆ than the prior experiment. PPPL estimates that it purchases from 2,400 to 4,700 pounds of SF₆ annually, and about 1,100 to 2,300 pounds are assumed to leak into the atmosphere annually. Some of the gas is contained in the equipment. During maintenance on equipment that requires removal of the SF₆ gas, PPPL uses SF₆ reclamation equipment to recover, clean, and reuse as much SF₆ gas as possible from the equipment.⁶⁶

Relative to total industrial non-combustion process emissions, SF₆ emissions from PPPL equipment are estimated to average about 0.018 MMtCO₂e annually from 1998 through 2005. Based on the range of gas leaks provided by PPPL, emissions may be as low as 0.012 MMtCO₂e or as high as 0.025 MMtCO₂e annually. For the purpose of this preliminary inventory, emissions for 1990 through 1997 were estimated to be about 0.325 MMtCO₂e per year based on an assumption that 30,000 pounds per year of SF₆ gas was released to the atmosphere annually. Based on the range of gas leaks provided by PPPL, emissions may be as low as 0.304 MMtCO₂e or as high as 0.39 MMtCO₂e annually for 1990 through 1997. Because of scaling effects, emissions for this category cannot be seen in Figure D2 after year 2000.

Nitric Acid Production

The manufacture of nitric acid (HNO₃) produces N₂O as a by-product, via the oxidation of ammonia. Nitric acid is a raw material used primarily to make synthetic commercial fertilizer. It is also a major component in the production of adipic acid (a feedstock for nylon) and explosives. Relatively small quantities of nitric acid are also employed for stainless steel pickling, metal etching, rocket propellants, and nuclear fuel processing.⁶⁷ The NJDEP provided nitric acid production data from its *Release and Pollution Prevention Reports*. Data were provided for EI DuPont Denemours & Co., Inc for 1994-2003, and for Saint Gobain Performance Plastics Corp. that produced nitric acid in 1998 only. Total production for each year was entered into the SGIT to calculate GHG emissions. The SGIT uses a default emission factor of 0.008 metric tons of N₂O emissions per metric ton of nitric acid produced based on a weighted-average calculated over the different types of emissions control technologies typically employed by nitric acid plants nationwide.⁶⁸

The NJDEP and BPU prepared a GHG emissions inventory for 1990. The documentation for this inventory provided 1990 production of nitric acid for one plant of 119,070 metric tons (131,250 short tons) and an emission factor of 0.0055 tons of N₂O per ton of nitric acid

⁶⁵ NJDEP and NJBPU, 1996, New Jersey Inventory of Greenhouse Gas Emissions, 1990, prepared by Cameron Johnson and Michael Aucott, Division of Science and Research, NJDEP, Trenton, NJ.

⁶⁶ E-mail from Jerry Levine, PPPL, to Mike Aucott, NJDEP, May 31, 2007.

⁶⁷ EIIP, Volume VIII: Chapter. 6. "Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes", August 2004.

⁶⁸ According to Chapter 6 of the EIIP guidance document, the nitric industry controls for oxides of nitrogen through two technologies: non-selective catalytic reduction (NSCR) and SCR. Only one of these technologies, NSCR, is effective at destroying N₂O emissions in the process of destroying oxides of nitrogen emissions. NSCR technology was widely installed in nitric acid plants built between 1971 and 1977. Due to high-energy costs and associated high gas temperatures, this technology has not been popular with modern plants. Only about 20% of the current plants have NSCR technology installed. All other plants have installed SCR technology. Since 80% of the current plants have SCR technology installed and 20% have NSCR technology, the weighted-average emission factor used in the SGIT is equal to $(0.0095 \times 0.80) + (0.002 \times 0.20) = 0.008$ metric tons N₂O per metric ton of nitric acid produced.

produced.⁶⁹ This information was used to estimate emissions for 1990, and a GWP of 310 was used to convert emissions to a MMtCO₂e basis.

Growth rates for New Jersey's manufacturing sector for 2006 through 2020 were used to forecast emissions (see Table D2). Emissions for 2004 and 2005 were held constant at 2003 levels (i.e., the last year for which production data were available). Relative to total industrial non-combustion process emissions, estimated emissions associated with nitric acid production are low (about 0.0003 MMtCO₂e in 1995 and 0.001 MMtCO₂e in 2020), and therefore, appear at the bottom of Figure D2 due to scaling effects.

Semiconductor Manufacture

The semiconductor industry uses fluorinated gases (PFCs [CF₄, C₂F₆, and C₃F₈]; HFC-23; and SF₆) in plasma etching and chemical vapor deposition processes. Emissions of SF₆ and HFCs from the manufacture of semiconductors have experienced declines since 2000). Emissions for New Jersey from 1990 to 2002 were estimated based on the default estimates provided in SGIT, which uses the ratio of the state-to-national value of semiconductor shipments to estimate the state's proportion of national emissions from the US EPA GHG inventory (US EPA 2007 *Inventory of US Greenhouse Gas Emissions and Sinks: 1990-2005*). The US Climate Action Report shows expected decreases in these emissions at the national level, and the same rate of decline is assumed for emissions in New Jersey. The decline in emissions in the future reflects expectations of future actions by the semiconductor industry to reduce these emissions. Relative to total industrial non-combustion process emissions, emissions associated with the manufacture of semiconductors are low (about 0.011 MMtCO₂e in 1990, 0.025 MMtCO₂e in 2005, and projected to return to 1990 levels of 0.011 MMtCO₂e in 2020), and therefore appear at the bottom of Figure D2 due to scaling effects.

Limestone Consumption

Limestone and dolomite are basic raw materials used by a wide variety of industries, including the construction, agriculture, chemical, glass manufacturing, and environmental pollution control industries, as well as in metallurgical industries such as magnesium production.⁷⁰ Historical state-specific data for New Jersey were not available from the United States Geological Survey (USGS); consequently, the default methods used in SGIT to allocate regional USGS data to New Jersey were used to calculate emissions. Growth rates for New Jersey's manufacturing sector for 2006 through 2020 were used to forecast emissions (see Table D2). Emissions for 2004 and 2005 were held constant at 2003 levels (i.e., the last year for which production data were available). Relative to total industrial non-combustion process emissions, estimated emissions associated with limestone and dolomite consumption are low (about 0.013 MMtCO₂e in 1995 and 0.004 MMtCO₂e in 2020), and therefore, cannot be seen in Figure D2 due to scaling effects. Note that SGIT did not provide any dolomite consumption data for any historical year for New Jersey.

⁶⁹ New Jersey Inventory of Greenhouse Gas Emissions 1990, prepared by NJDEP and New Jersey BPU, 1996, Chapter 2, page 20.

⁷⁰ In accordance with EIIP Chapter 6 methods, emissions associated with the following uses of limestone and dolomite are not included in this category: (1) crushed limestone consumed for road construction or similar uses (because these uses do not result in CO₂ emissions), (2) limestone used for agricultural purposes (which is counted under the methods for the agricultural sector), and (3) limestone used in cement production (which is counted in the methods for cement production).

Soda Ash Consumption

Commercial soda ash (sodium carbonate) is used in many consumer products such as glass, soap and detergents, paper, textiles, and food. Carbon dioxide is also released when soda ash is consumed. SGIT estimates historical emissions based on the state's population and national per capita emissions from the US EPA national GHG inventory. According to the USGS, this industry is expected to grow at an annual rate of 0.5% from 2004 through 2009 for the US as a whole. Information on growth trends for years later than 2009 was not available; therefore the same 0.5% annual growth rate was applied for estimating emissions to 2020. Relative to total industrial non-combustion process emissions, emissions associated with soda ash consumption are low (about 0.084 MMtCO₂e in 1990 and 0.081 MMtCO₂e in 2020), and therefore appear at the bottom of Figure D2 due to scaling effects.

Key Uncertainties

Key sources of uncertainty underlying the estimates above are as follows:

- Since emissions from industrial processes are determined by the level of production and the production processes of a few key industries—and in some cases, a few key plants—there is relatively high uncertainty regarding future emissions from the industrial processes category as a whole. Future emissions depend on the competitiveness of New Jersey manufacturers in these industries, and the specific nature of the production processes used in New Jersey.
- The projected largest source of future industrial emissions, HFCs and PFCs used in cooling applications, is subject to several uncertainties as well. First, historical emissions are based on national estimates; New Jersey-specific estimates are currently unavailable. In addition, emissions through 2020 and beyond will be driven by future choices regarding mobile and stationary air conditioning technologies and the use of refrigerants in commercial applications, for which several options currently exist.
- One plant in New Jersey has produced nitric acid annually since 1994. A second plant reported the production of nitric acid in 1998 only. The degree to which the one plant has adopted or will adopt measures to reduce N₂O emissions is not specifically known. In addition, use of the employment growth rate to forecast emissions for this one plant may be highly uncertain. Consequently, there is uncertainty in the estimation of both historical and future emissions from nitric acid production.
- Greenhouse gases are emitted from several additional industrial processes that are not covered in the EIIP guidance documents, due in part to a lack of sufficient state data on non-energy uses of fossil fuels for these industrial processes. These sources include:
 - Titanium Dioxide Production (CO₂);
 - Phosphoric Acid Production (CO₂);
 - CO₂ Consumption (CO₂);
 - Ferroalloy Production (CO₂);
 - Petrochemical Production (CH₄); and
 - Silicon Carbide Production (CH₄).

The CO₂ emissions from the above CO₂ sources (other than CO₂ consumption and phosphoric acid production) result from the non-energy use of fossil fuels. Although the US EPA estimates emissions for these industries on a national basis, the US EPA has not developed methods for

estimating the emissions at the state level due to data limitations. If state-level data on non-energy uses of fuels become available, future work should include an assessment of emissions for these other categories.

Certain emission sources in New Jersey are required through the Emissions Statement Program to submit to the NJDEP annually emissions of volatile organic compounds, oxides of nitrogen, carbon monoxide, and other pollutants including CO₂ and CH₄. The emission sources must certify that the data they submit are accurate. Thus, this information could serve as a good starting point for building a bottom-up GHG inventory for industrial non-energy uses of fuels and processing of raw materials that contribute to GHG emissions. Some wastewater treatment sources also report under the Emission Statement Program. A preliminary review of this information indicated that it will be difficult and time-intensive to reconcile the top-down inventory approach used in this preliminary analysis with the Emissions Statement Data. A major contributor to this difficulty is the fact that only a subset of sources of releases of GHGs are currently required to report. The NJDEP anticipates using the Emission Statement Data in the future as it improves upon the methods and data sources used to construct its GHG inventory.

Appendix E. Fossil Fuel Extraction and Distribution Industry

Overview

The inventory for this subsector of the Energy Supply sector includes only methane (CH₄) emissions associated with the transmission and distribution (T&D) of natural gas in New Jersey. There is no oil or natural gas production or processing or coal mining in New Jersey. In 2005, emissions from natural gas T&D account for an estimated 2.0 million metric tons (MMt) of CO₂ equivalent (CO₂e) of total gross greenhouse gas (GHG) emissions in New Jersey, and are estimated to increase slightly to about 2.1 MMtCO₂e by 2020.

The following discusses the data sources, methods, assumptions, and results used to construct the inventory and reference case projections for this sector. The reference case projections assume business-as-usual practices.

Natural Gas T&D Emissions and Reference Case Projections (Business-as-Usual)

Methane emissions for 1990 through 2005 were estimated using the United States Environmental Protection Agency's (US EPA) State Greenhouse Gas Inventory Tool (SGIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for natural gas T&D.⁷¹ Table E1 provides an overview of the required data, data sources, and the approach to projecting future emissions. The activity data were entered into the SGIT to calculate emissions for 1990 through 2005. Methane emissions were calculated by multiplying emissions-related activity levels (e.g., miles of pipeline) by aggregate emission factors.

The annual activity data for miles of T&D pipeline and number of service connections to distribution pipeline were obtained from databases provided by the Office of Pipeline Safety (OPS).⁷² For the distribution system, annual CH₄ emissions were estimated using OPS' pipeline mileage and the SGIT emission factors for (1) distribution pipeline constructed of cast iron, unprotected steel, protected steel, and plastic, and (2) the number of protected and unprotected service connections. For the transmission system, the SGIT methods use total miles of pipeline as the basis for calculating CH₄ emissions; separate emission factors are not provided for pipeline constructed of different materials.

The SGIT methods also include emission factors for estimating CH₄ emissions associated with leaks from gas transmission compressor stations, gas storage compressor stations, and liquefied natural gas (LNG) storage compressor stations. Information on the type and number of compressor stations was not readily available for New Jersey. Therefore, the default factors in SGIT for estimating the number of gas transmission compressor stations and gas storage compressor stations from the miles of transmission pipeline for each year were used. The SGIT assumes 0.006 gas transmission compressor station per mile of transmission pipeline and 0.0015 gas storage compressor station per mile of transmission pipeline. Note that the SGIT

⁷¹ Methane emissions were calculated using SGIT, with reference to Emission Inventory Improvement Program, Volume VIII: Chapter. 5. "Methods for Estimating Methane Emissions from Natural Gas and Oil Systems", March 2005.

⁷² US Office of Pipeline Safety, Distribution and Transmission Annuals Data for 1990-2005, <http://ops.dot.gov/stats/DT98.htm>.

does not provide default methods for estimating the number of LNG storage compressor stations in a state; therefore, emissions were not estimated for LNG storage compressor stations.

Table E1 shows the compound annual average growth rates applied to forecast emissions associated with natural gas T&D from 2006 through 2020. These annual growth assumptions are based on average annual growth in natural gas consumption in New Jersey calculated from the US Department of Energy (DOE), Energy Information Administration’s (EIA) AEO2006 regional forecast for the EIA’s Mid-Atlantic region.⁷³

Table E1. Approach to Estimating Historical and Future Methane Emissions from Natural Gas Transmission and Distribution

	Approach to Estimating Historical Emissions		Approach to Estimating Projections
Activity	Required Data for SGIT	Data Source	Projection Assumptions
Natural Gas Transmission	Miles of transmission pipeline	OPS	Based on average annual growth in natural gas consumption in New Jersey calculated from AEO2006 regional forecast for Energy Information Administration’s Mid-Atlantic region: 0.38% for 2005-2010, 0.45% for 2010-2015, and 0.35% for 2015-2020.
	Number of gas transmission compressor stations	US EPA EIIP default assumptions	
	Number of gas storage compressor stations	US EPA EIIP default assumptions	
	Number of LNG storage compressor stations	No data available	
Natural Gas Distribution	Miles of distribution pipeline	OPS	
	Total number of services	OPS	
	Number of unprotected steel services	OPS	
	Number of protected steel services	OPS	

Results


Figure E1 displays the estimated GHG emissions associated with natural gas T&D system in New Jersey from 1990 to 2005, with projections to 2020. Figure E1 was developed from the emissions data in Table E1. Table E2 shows the relative contributions of emissions associated with the distribution and transmission systems to total subsector emissions.

Emissions associated with this sector are estimated to be about 2.1 MMtCO₂e in 1990, 2.0 MMtCO₂e in 2005, and 2.1 MMtCO₂e in 2020. As shown in Figure E1, CH₄ emissions associated with New Jersey’s distribution system declined from 1990 through 2000. This decline in emissions is associated with natural gas companies in New Jersey replacing cast iron and unprotected steel distribution pipe with protected steel and plastic pipe. Gas companies also replaced unprotected steel service connections with protected steel and plastic service connections that helped reduce emissions during this ten-year period. The increase in distribution emissions after 1995 is associated with the expansion of the distribution pipeline

⁷³ US DOE, EIA AEO2006 with Projections to 2030, <http://www.eia.doe.gov/oiaf/aeo/index.html>.

system to accommodate growth using cathodically protected and coated or plastic pipeline and service connections. The increase in emissions associated with the transmission system from 2000 to 2005 reflects an increase in the on-shore transmission pipeline mileage over this period. According to the OPS data, on-shore transmission pipeline mileage increased from 404 miles in 2000 to 1,427 miles in 2005.⁷⁴

Figure E1. Methane Emissions and Projections from the Fossil Fuel Industry



Source: Results in graph based on approach described in text.

Table E1. Emissions Inventory and Reference Case Projections (MMtCO₂e)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Transmission (CH ₄)	0.07	0.09	0.07	0.24	0.24	0.25	0.25
Distribution (CH ₄)	1.99	1.87	1.80	1.78	1.82	1.86	1.89
Total	2.06	1.95	1.87	2.02	2.06	2.10	2.14

Source: Results in table based on approach described in text.

⁷⁴ Note that the OPS data indicate that New Jersey does not have any on-shore or off-shore gathering or off-shore transmission pipeline.

Table E2. Proportions of Total Subsector Emissions by Type and Source (%)

Fuel Type	1990	1995	2000	2005	2010	2015	2020
Transmission (CH ₄)	3%	4%	4%	12%	12%	12%	12%
Distribution (CH ₄)	97%	96%	96%	88%	88%	88%	88%

Source: Results in table based on approach described in text.

Key Uncertainties

The main uncertainties are associated with the reference case projection assumptions. For this preliminary forecast, it was assumed that emissions would increase at the rate of natural gas consumption that the EIA projected in its AEO2006 regional forecast for the EIA's Mid-Atlantic region. Market factors (e.g., price of natural gas relative to other available energy sources) could have a significant impact on the growth for this sector. In addition, neither potential future application of improvements to pipeline technologies that can yield emission reductions nor the potential effect of demand-side management programs in reducing gas consumption have been accounted for in the emissions projections shown here.

Future improvements to the estimates for the inventory should include the collection of activity data from gas companies to (1) verify the OPS data used for the miles of T&D pipeline and distribution service connections, (2) replace the SGIT defaults for estimating the number of gas transmission compressor stations and gas storage compressor stations, and (3) estimate emissions associated with LNG storage compressor stations if it is determined that these stations exist in New Jersey.

Appendix F. Agriculture

Overview

The emissions discussed in this appendix refer to non-energy methane (CH₄) and nitrous oxide (N₂O) emissions from enteric fermentation, manure management, and agricultural soils. Emissions and sinks of carbon in agricultural soils are also covered. Energy emissions (combustion of fossil fuels in agricultural equipment) are included in the residential, commercial, and industrial (RCI) sector estimates (see Appendix B).

There are two livestock sources of greenhouse gas (GHG) emissions: enteric fermentation and manure management. Methane emissions from enteric fermentation are the result of normal digestive processes in ruminant and non-ruminant livestock. Microbes in the animal digestive system break down food and emit CH₄ as a by-product. More CH₄ is produced in ruminant livestock because of digestive activity in the large fore-stomach. Methane and N₂O emissions from the storage and treatment of livestock manure (e.g., in compost piles or anaerobic treatment lagoons) occur as a result of manure decomposition. The environmental conditions of decomposition drive the relative magnitude of emissions. In general, the more anaerobic the conditions are, the more CH₄ is produced because decomposition is aided by CH₄ producing bacteria that thrive in oxygen-limited conditions. Under aerobic conditions, N₂O emissions are dominant. Emissions estimates from manure management are based on manure that is stored and treated on livestock operations. Emissions from manure that is applied to agricultural soils as an amendment or deposited directly to pasture and grazing land by grazing animals are accounted for in the agricultural soils emissions.

The management of agricultural soils can result in N₂O emissions and net fluxes of carbon dioxide (CO₂) causing emissions or sinks. In general, soil amendments that add nitrogen to soils can also result in N₂O emissions. Nitrogen additions drive underlying soil nitrification and denitrification cycles, which produce N₂O as a by-product. The emissions estimation methodologies used in this inventory account for several sources of N₂O emissions from agricultural soils, including decomposition of crop residues, synthetic and organic fertilizer application, manure application, sewage sludge, nitrogen fixation, and histosols (high organic soils, such as wetlands or peatlands) cultivation. Both direct and indirect emissions of N₂O occur from the application of manure, fertilizer, and sewage sludge to agricultural soils. Direct emissions occur at the site of application and indirect emissions occur when nitrogen volatilizes or leaches to groundwater or in surface runoff and is transported off-site before entering the nitrification/denitrification cycle. Methane and N₂O emissions also result when crop residues are burned and during rice cultivation; however, open burning of crops is prohibited in New Jersey due to air quality concerns and rice is not grown in New Jersey.

The net flux of CO₂ in agricultural soils depends on the balance of carbon losses from management practices and gains from organic matter inputs to the soil. Carbon dioxide is absorbed by plants through photosynthesis and ultimately becomes the carbon source for organic matter inputs to agricultural soils. When inputs are greater than losses, the soil accumulates carbon and there is a net sink of CO₂ into agricultural soils. In addition, soil disturbance from the cultivation of histosols releases large stores of carbon from the soil to the atmosphere. Finally, the practice of adding limestone and dolomite to agricultural soils results in CO₂ emissions.

The following discusses the data sources, methods, assumptions, and results used to construct the inventory and reference case projections for this sector. The reference case projections assume business-as-usual practices.

Emissions and Reference Case Projections (Business-as-Usual)

Methane and Nitrous Oxide

GHG emissions for 1990 through 2005 were estimated using the United States Environmental Protection Agency's (US EPA) State Greenhouse Gas Inventory Tool (SGIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for the sector.⁷⁵ In general, the SGIT methodology applies emission factors developed for the US to activity data for the agriculture sector. Activity data include livestock population statistics, crop production statistics, amounts of fertilizer applied to crops, and trends in manure management practices. This methodology is based on international guidelines developed by sector experts for preparing GHG emissions inventories.⁷⁶

Defaults in SGIT for New Jersey livestock population and crop production from 1990 to 2005 come from the USDA National Agriculture Statistical Service (NASS).⁷⁷ These default data were compared to data from New Jersey Department of Agriculture Annual Reports⁷⁸ and replaced with the local data where differences were found. The default SGIT manure management system assumptions for each livestock category were used for this inventory. SGIT data on fertilizer usage came from *Commercial Fertilizers*, a report from the Fertilizer Institute. Activity data for fertilizer includes all potential uses in addition to agriculture, such as residential and commercial (e.g., golf courses). The estimates are reported in the agriculture sector but they represent emissions occurring on other land uses.

Data were not available to estimate nitrogen released by the cultivation of histosols (i.e., the number of acres of high organic content soils). Given that cultivation of organic soils is a source of CO₂ emissions in New Jersey (see below), N₂O emissions are also probably occurring.

Emissions from enteric fermentation and manure management were projected based on forecasted animal populations. Dairy cattle forecasts were based on state-level projections of dairy cows from the Food and Agricultural Policy Research Institute (FAPRI).⁷⁹ Projections for all other livestock categories, except swine, were estimated based on linear forecasts of the historical 1990-2005 populations. The swine population showed a sharp decline in the late 1990s, and linear projection of the 1990-2005 populations result in negative populations before

⁷⁵ GHG emissions were calculated using SGIT, with reference to EIIP, Volume VIII: Chapter 8. "Methods for Estimating Greenhouse Gas Emissions from Livestock Manure Management", August 2004; Chapter 10. "Methods for Estimating Greenhouse Gas Emissions from Agricultural Soil Management", August 2004; and Chapter 11. "Methods for Estimating Greenhouse Gas Emissions from Field Burning of Agricultural Residues", August 2004.

⁷⁶ Revised 1996 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National Greenhouse Gas Inventories, published by the National Greenhouse Gas Inventory Program of the IPCC, available at (<http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.htm>); and Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories, published in 2000 by the National Greenhouse Gas Inventory Program of the IPCC, available at: (<http://www.ipcc-nggip.iges.or.jp/public/gp/english/>).

⁷⁷ USDA, NASS (http://www.nass.usda.gov/Statistics_by_State/New_Jersey/index.asp).

⁷⁸ Annual Reports, New Jersey Department of Agriculture, <http://www.state.nj.us/agriculture/pub/general.html>.

⁷⁹ FAPRI Agricultural Outlook 2006, Food and Agricultural Policy Research Institute, <http://www.fapri.iastate.edu/outlook2006>.

2020. Therefore, swine populations were projected based on the 2000-2005 historical data. Table F1 shows the 2005-2020 annual growth rates estimated for each category.

Table F1. Growth Rates Applied for the Enteric Fermentation and Manure Management Categories

Livestock Category	2005-2020 Annual Growth
Dairy Cattle	-2.6%
Beef Cattle	-1.2%
Swine	-13.1%
Sheep	-5.2%
Goats	1.5%
Horses	-3.4%
Turkeys	-2.7%
Layers	0.6%

Projections for agricultural soils categories were based on linear extrapolation of the 1990-2005 historical data. Table F2 shows the 2005-2020 annual growth rates estimated for each category.

Table F2. Growth Rates Applied for the Agricultural Sector

Agricultural Category	Growth Rate
Direct	
Fertilizers	0.9%
Crop Residues	-3.5%
Nitrogen-Fixing Crops	-2.4%
Livestock	-4.6%
Indirect	
Fertilizers	0.9%
Livestock	-5.3%
Leaching/Runoff - Fertilizer	0.9%
Leaching/Runoff - Livestock	-5.3%

Soil Carbon

Net carbon fluxes from agricultural soils have been estimated by researchers at the Natural Resources Ecology Laboratory at Colorado State University and are reported in the US Inventory of Greenhouse Gas Emissions and Sinks⁸⁰ and the US Agriculture and Forestry Greenhouse Gas Inventory. The estimates are based on the Intergovernmental Panel on Climate Change (IPCC) methodology for soil carbon adapted to conditions in the US. Preliminary state-level estimates of CO₂ fluxes from mineral soils and emissions from the cultivation of organic soils were reported in the US Agriculture and Forestry Greenhouse Gas Inventory.⁷ Currently, these are the best available data at the state-level for this category. The inventory did not report state-level estimates of CO₂ emissions from limestone and dolomite applications; hence, this source is not included in this inventory at present.

Carbon dioxide fluxes resulting from specific management practices were reported. These practices include: conversions of cropland resulting in either higher or lower soil carbon levels;

⁸⁰ US Inventory of Greenhouse Gas Emissions and Sinks: 1990-2004 (and earlier editions), US Environmental Protection Agency, Report # 430-R-06-002, April 2006. Available at: <http://www.epa.gov/climatechange/emissions/usinventoryreport.html>.

additions of manure; participation in the Federal Conservation Reserve Program (CRP); and cultivation of organic soils (with high organic carbon levels). For New Jersey, Table F3 shows a summary of the latest estimates available from the USDA, which are for 1997.⁸¹ These data show that changes in agricultural practices are estimated to result in a net sink of 0.07 million metric tons (MMt) of CO₂ equivalent (CO₂e) per year (yr) in New Jersey. Since data are not yet available from USDA to make a determination of whether the emissions are increasing or decreasing, a sink of 0.07 MMtCO₂e/yr is assumed to remain constant.

Table F3. GHG Emissions from Soil Carbon Changes Due to Cultivation Practices (MMtCO₂e)

Changes in cropland			Changes in Hayland				Other			Total ⁴
Plowout of grassland to annual cropland ¹	Cropland management	Other cropland ²	Cropland converted to hayland ³	Hayland management	Cropland converted to grazing land ³	Grazing land management	CRP	Manure application	Cultivation of organic soils	Net soil carbon emissions
0.11	(0.04)	0.00	(0.11)	0.00	(0.04)	0.00	0.00	(0.04)	0.04	(0.07)

Based on USDA 1997 estimates. Parentheses indicate net sequestration.

¹ Losses from annual cropping systems due to plow-out of pastures, rangeland, hayland, set-aside lands, and perennial/horticultural cropland (annual cropping systems on mineral soils, e.g., corn, soybean, cotton, and wheat).

² Perennial/horticultural cropland and rice cultivation.

³ Gains in soil carbon sequestration due to land conversions from annual cropland into hay or grazing land.

⁴ Total does not include change in soil organic carbon storage on federal lands, including those that were previously under private ownership, and does not include carbon storage due to sewage sludge applications.

Results

Fertilizers are the largest contributor to agricultural greenhouse gas emissions in New Jersey. This category includes direct and indirect emissions of N₂O from synthetic and non-manure organic fertilizers. In 1990, fertilizers accounted for 36% (0.22 MMtCO₂e) of total agricultural emissions. Fertilizer emissions increased slightly to 0.23 MMtCO₂e (47% of total agricultural emissions) in 2005. While other agricultural emissions categories are expected to decline, fertilizer emissions are estimated to increase to 0.27 MMtCO₂e in 2020. The high percentage of agricultural emissions associated with fertilizers and the upward trend of this category is likely due to the fact that a high percentage of fertilizer usage in New Jersey is for non-farm applications. According to the 2006 New Jersey Department of Agriculture Annual Report⁸², about 48% of total mixed fertilizers and materials was used for non-farm applications.

The agricultural soils-livestock category, which includes direct and indirect N₂O emissions from manure applications and direct deposition of manure on agricultural soils, accounted for 25% (0.15 MMtCO₂e) of total agricultural emissions in 1990. Agricultural soils-livestock emissions decreased to 0.11 MMtCO₂e (21% of total agricultural emissions) in 2005. Emissions from this

⁸¹ US Agriculture and Forestry Greenhouse Gas Inventory: 1990-2001. Global Change Program Office, Office of the Chief Economist, US Department of Agriculture. Technical Bulletin No. 1907, 164 pp. March 2004. http://www.usda.gov/oce/global_change/gg_inventory.htm; the data are in appendix B table B-11. The table contains two separate IPCC categories: “carbon stock fluxes in mineral soils” and “cultivation of organic soils.” The latter is shown in the second to last column of Table F1. The sum of the first nine columns is equivalent to the mineral soils category.

⁸² “Mixed Fertilizers, Fiscal Year Ending June 30, 2006”, Page 93, New Jersey Agriculture 2006 Annual Report, <http://www.state.nj.us/agriculture/pdf/06AnnualReport.pdf>.

category are expected to continue declining to 0.05 MMtCO₂e in 2020, due to the projected decline in livestock populations.

The agricultural soils-crops category decreased between 1990 to 2005, with 1990 emissions accounting for 13% (0.08 MMtCO₂e) of total agricultural emissions and 2005 emissions estimated to be 11% (0.05 MMtCO₂e) of total agricultural emissions. Emissions from this category are estimated to be 0.04 MMtCO₂e in 2020.

In 1990, enteric fermentation accounted for about 21% (0.13 MMtCO₂e) of total agricultural emissions. Enteric fermentation emissions decreased to 0.08 MMtCO₂e (16% of total agricultural emissions) in 2005 due primarily to the decline in beef and dairy cattle populations between 1990 and 2005. Cattle populations are projected to continue declining through 2020, and enteric fermentation emissions are estimated to be 0.06 MMtCO₂e in 2020.


The manure management category accounted for only 6% (0.04 MMtCO₂e) of total agricultural emissions in 1990 and decreased to 5% (0.03 MMtCO₂e) in 2005. Manure management emissions are estimated to continue declining to 0.02 MMtCO₂e of total agricultural emissions in 2020, due to declines in livestock populations.

The only standard IPCC source categories missing from this report are CO₂ emissions from limestone and dolomite application and N₂O emissions from the cultivation of histosols. Estimates for limestone and dolomite application in New Jersey were not available; however, the USDA's national estimate for soil liming is about 9 MMtCO₂e/yr.⁸³

GHG emissions from the agriculture sector in New Jersey are summarized in Figure F1 and Table F4.

⁸³ US Agriculture and Forestry Greenhouse Gas Inventory: 1990-2001. Global Change Program Office, Office of the Chief Economist, US Department of Agriculture. Technical Bulletin No. 1907. 164 pp. March 2004.

Figure F1. Gross GHG Emissions from Agriculture



Notes: Ag Soils – Crops category includes: incorporation of crop residues and nitrogen fixing crops (no cultivation of histosols estimated); Ag Soils – Fertilizers category includes farm and non-farm utilization of fertilizers.

Table F4. Net GHG Emissions from Agriculture (MMtCO₂e)

Source	1990	1995	2000	2005	2010	2015	2020
Enteric Fermentation	0.128	0.116	0.090	0.080	0.070	0.064	0.059
Manure Management	0.038	0.037	0.031	0.026	0.027	0.025	0.022
Ag Soils-Fertilizers	0.219	0.184	0.231	0.232	0.247	0.256	0.265
Ag Soils-Crops	0.078	0.066	0.080	0.053	0.052	0.044	0.035
Ag Soils-Livestock	0.152	0.135	0.117	0.105	0.086	0.068	0.050
Agricultural Burning	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Total Gross Emissions	0.615	0.537	0.548	0.497	0.482	0.457	0.432
Net Soil Carbon Emissions	-0.07	-0.07	-0.07	-0.07	-0.07	-0.07	-0.07
Total Net Emissions	0.545	0.467	0.478	0.427	0.412	0.387	0.362

Key Uncertainties

Emissions from enteric fermentation and manure management are dependent on the estimates of animal populations and the various factors used to estimate emissions for each animal type and manure management system (i.e., emission factors which are derived from several variables including manure production levels, volatile solids content, and CH₄ formation potential). Each of these factors has some level of uncertainty. Also, animal populations fluctuate throughout the year, and thus using point estimates introduces uncertainty into the average annual estimates of these populations. The largest contributors to uncertainty in emissions from manure management are the emission factors, which are derived from limited data sets.

As mentioned above, for emissions associated with changes in agricultural soil carbon levels, the only data currently available are for 1997. When newer data are released by the USDA, these should be reviewed to represent current conditions as well as to assess trends. In particular, given the potential for some CRP acreage to retire and possibly return to active cultivation prior to

2020, the emissions could be appreciably affected. As mentioned above, emission estimates for soil liming and cultivation of histosols have not been developed for New Jersey.

Another contributor to the uncertainty in the emission estimates is the projection assumptions. The growth rates for most source categories are based on the assumption that the average annual rate of change in future year emissions will follow the historical average annual rate of change from 1990 through 2005.

Appendix G. Waste Management

Overview

Greenhouse gas (GHG) emissions from waste management include:

- Solid waste management – methane (CH₄) emissions from municipal and industrial solid waste landfills, accounting for CH₄ that is flared or captured for energy production (this includes both open and closed landfills);
- Solid waste combustion – CH₄, carbon dioxide (CO₂), and nitrous oxide (N₂O) emissions from the combustion of solid waste in incinerators or waste to energy plants or during open burning; and
- Wastewater management – CH₄ and N₂O from municipal wastewater and CH₄ from industrial wastewater (WW) treatment facilities.

The following discusses the data sources, methods, assumptions, and results used to construct the inventory and reference case projections for this sector. The reference case projections assume business-as-usual practices.

Inventory and Reference Case Projections (Business-as-Usual)

Solid Waste Management

For solid waste management, emissions were calculated using the United States Environmental Protection Agency's (US EPA) AP-42 equation,⁸⁴ which is used by EPA's Landfill Gas Emissions Model (LandGEM). The New Jersey Department of Environmental Protection (NJDEP) provided input data to estimate annual waste emplacement and CH₄ emissions for each landfill.

The NJDEP provided additional information for landfills to identify them as being uncontrolled, flared, passively vented, or with landfill gas to energy (LFGTE) projects. Small uncontrolled landfills in some counties were combined for the purposes of emissions modeling. There are 89 large landfill sites representing both open and closed landfills. Fifteen of these sites collect landfill gas for use in an LFGTE plant. Another 17 sites collect and flare landfill gas. The remaining 67 sites are uncontrolled or passively vented. Table G1 provides a listing of sites that are controlled.

There are some additional CH₄ emissions from the approximately 300 landfills that are too small and old to be listed individually. NJDEP extrapolated a best-fitting exponential trend line of a plot of landfill volume versus estimated methane emissions data, which suggested that the total methane emissions from these landfills is no more than 6,000 short tons/year.

⁸⁴ As per EPA's AP-42 Section on Municipal Solid Waste Landfills:
<http://www.epa.gov/ttn/chief/ap42/ch02/final/c02s04.pdf>.

Table G1. Controlled Solid Waste Landfills

LFGTE Landfills		Flared Landfills
Name	LFG Use	Name
Cape May County	Heat/Steam	Big Hill
HMDC 1-A	Heat/Steam	Burlington County
HMDC 1-D	Heat/Steam	Buzby Bros.
BCUA Kingsland	Electricity	Combe Fill North
Edgeboro	Electricity	Combe Fill South
Edison Disposal Area	Electricity	Cumberland County
Hamms Landfill	Electricity	Florence
HMDC 1-E, Balefill 1C	Electricity	Gems
ILR	Electricity	Gloucester County
Kinsley's	Electricity	Kin-Buc
Middlesex County LF	Electricity	Kramer
Monmouth County Rec Ctr PH II	Electricity	L & D Mt. Holly
Monmouth County Rec Ctr PH I	Electricity	Parklands
Ocean County	Electricity	Pennsauken
Sussex County 1-E	Electricity	Pineland Park
		Salem County Landfill
		SLF Inc

Annual emissions from municipal solid waste (MSW) landfills were estimated and categorized as follows: (1) uncontrolled landfills; (2) landfills with a landfill gas collection system and LFGTE plant; and (3) landfills with landfill gas collection and a flare. For the controlled landfills, it was assumed that the overall methane collection and control efficiency is 75%.⁸⁵ Of the methane not captured by a landfill gas collection system, it is further assumed that 10% is oxidized before being emitted to the atmosphere (consistent with the EPA State Greenhouse Gas Inventory Tool [SGIT] default assumption).

For the period from 1990 through 1995, many of the LFTGE and some of the flaring operations were not yet in place at the New Jersey landfills. To account for this, NJDEP provided estimates of the 1990 and 1995 emissions from in-state landfills, calculated with an earlier, simpler method than the EPA exponential formula. This resulted in landfill emission values of 9.8 MMtCO₂e in 1990 and 6.6 MMtCO₂e in 1995.⁸⁶ Values for the interim years were calculated via interpolation.

In addition to the in-state landfills, New Jersey exports a significant amount of waste for disposal out-of-state. NJDEP provided data on the amount of waste disposed of in-state and the amount disposed of out-of-state from 1985 through 2004.⁸⁷ NJDEP separately provided information on the amount of the waste disposed of in-state that was incinerated in 2003.⁸⁸ To estimate the

⁸⁵ As per EPA's AP-42 Section on Municipal Solid Waste Landfills:
<http://www.epa.gov/ttn/chief/ap42/ch02/final/c02s04.pdf>.

⁸⁶ Based on NJDEP calculations, personal communication, Michael Aucott.

⁸⁷ "New Jersey Solid Waste Database Trends Analysis: 1985 through 2004 (millions of tons per year)," http://www.state.nj.us/dep/dshw/recycling/05_trends.pdf.

⁸⁸ Michael Aucott, New Jersey Department of Environmental Protection, spreadsheet MSW_incin_calcs.xls, provided to Maureen Mullen, CCS, August 9, 2007.

amount of in-state waste incinerated in each remaining year from 1990 through 2004, the ratio of the tons of waste incinerated in-state in 2003 to the total tons of waste disposed of in-state in 2003 was multiplied by the total amount of waste disposed of in-state in the selected year. The CO₂ emissions from the active in-state landfills for each year from 1990 through 2004 were then totaled. To estimate the emissions from the waste generated in New Jersey, but exported out-of-state, these emissions from active in-state landfills were then multiplied by the ratio of the tons of waste disposed of out-of-state in a given year to the tons of waste landfilled in-state in that same year (total tons of waste disposed of in-state minus the calculated tons of waste incinerated).

Reference case projections of emissions from landfills were based on the same equations used to calculate the base inventory, with waste continuing to be emplaced at the landfills that are currently active (as of 2006). The ratio of the tons of waste from New Jersey landfilled out-of-state to the tons of waste landfilled in-state was assumed to remain at the 2004 level (the latest year for which these data were available) of 1.15 tons of waste landfilled out-of-state for every ton of waste landfilled in-state through the year 2020. The calculation of emissions from this waste disposed of out-of-state for the reference case projections was performed in the same manner as the calculation of emissions from out-of-state waste disposal used in the base inventory.

The SGIT default was used for industrial solid waste landfills. This default is based on national data indicating that industrial landfilled waste is emplaced at approximately 7% of the rate of MSW emplacement. We assumed that this additional industrial waste emplacement occurs beyond that already addressed in the emplacement rates for MSW sites described above. Due to a lack of data, no controls were assumed for industrial waste landfills. Emissions from industrial landfills from 2007 through 2020 were calculated using the same rate of 7% of the uncontrolled MSW emissions in each of the reference case projection years.

Solid Waste Combustion

NJDEP provided throughput data for the five municipal waste combustion facilities currently operating in New Jersey (Camden, Essex, Gloucester, Union, and Warren Counties).⁸⁹ Emissions from these waste combustion facilities are captured under the electricity production sector as these are all waste-to-energy facilities. Therefore, emissions from solid waste combustion are listed as 0 for the waste sector.

No readily-available data were identified to estimate emissions associated with residential open burning of solid waste. It is anticipated that the contribution from these practices would be negligible since open burning of waste is not permitted in New Jersey.

Wastewater Management

GHG emissions from municipal wastewater treatment were also estimated. Emissions are calculated in EPA's SGIT based on state population, assumed biochemical oxygen demand (BOD) and protein consumption per capita, and emission factors for N₂O. NJDEP provided data on CH₄ emissions in 2003 from 12 large New Jersey wastewater facilities that report to the NJDEP pursuant to its Emissions Statement Program. The NJDEP estimated that the flows from

⁸⁹ Michael Aucott, New Jersey Department of Environmental Protection.

these 12 facilities accounted for 65.8% of the total municipal wastewater flow in the State. The total CH₄ emissions from the 12 major facilities were divided by 65.8% to estimate the total CH₄ emissions from wastewater within the State. The resulting CH₄ wastewater emissions for 2003 were then adjusted to the other historical years from 1990 to 2005 in the same ratio as the historical year's SGIT-based N₂O emissions to the SGIT-based 2003 N₂O emissions. The key SGIT default values are shown in Table G2 below. The value of 85% of New Jersey residents not on septic differs from the SGIT default of 75%, based on information provided by NJDEP.⁹⁰ Wastewater emissions were projected from 2006 through 2020 based on the 1990-2005 historical growth rate (1.2%/yr).

Table G2. SGIT Key Default Values for Municipal Wastewater Treatment

Variable	Value
BOD	0.065 kg /day-person
Amount of BOD anaerobically treated	16.25%
CH ₄ emission factor	0.6 kg/kg BOD
New Jersey residents not on septic	85%
Water treatment N ₂ O emission factor	4.0 g N ₂ O/person-yr
Biosolids emission factor	0.01 kg N ₂ O-N/kg sewage-N
Source: U.S. EPA State Inventory Tool – Wastewater Module; methodology and factors taken from U.S. EPA, Emission Inventory Improvement Program, Volume 8, Chapter 12, October 1999: www.epa.gov/ttn/chieff/eiip/techreport/volume08/ .	


SGIT provides default assumptions and emission factors for industrial wastewater emissions from three industrial sectors: Fruit & Vegetables, Red Meat & Poultry, and Pulp & Paper. Data for wastewater from these industrial sectors could not be identified; therefore emissions for this category were not estimated.

Figure G1 and Table G3 show the emission estimates for the waste management sector. Overall, the sector accounts for 6.8 MMtCO₂e in 2005. By 2020, emissions are expected to decrease to 6.0 MMtCO₂e/yr. The decline in emissions is driven by the in-state solid waste landfill sector. In 2005, about 69% of the waste management sector emissions were contributed by in-state solid waste landfills. By 2020, in-state landfills are expected to contribute about 59% of the waste management sector emissions. Much of this decline is due to the continued degradation of the relatively large quantities of waste that were deposited in New Jersey landfills from out-of-state sources up until the late 1980s. As waste ages and decomposition proceeds, CH₄ emissions decline. In contrast, the share of waste management emissions from New Jersey waste landfilled out-of-state increases from 23% in 2005 to 30% in 2020.

In 2005, municipal wastewater contributed 8% of total waste management emissions. By 2020, the contribution from this sector is expected to be 11%. Note that N₂O emissions from municipal wastewater estimates are based on the default parameters listed in Table G1 above and might not adequately account for existing controls or management practices (e.g., anaerobic digesters served by a flare or other combustion devices).

⁹⁰ Michael Aucott, New Jersey Department of Environmental Protection.

Figure G1. New Jersey GHG Emissions from Waste Management



Notes: LF – landfill; WW – wastewater; LFGTE – landfill gas to energy.

Table G3. New Jersey GHG Emissions from Waste Management (MMtCO₂e)

Source	1990	1995	2000	2005	2010	2015	2020
Uncontrolled LFs	3.18	2.73	2.41	2.19	2.00	1.85	1.73
Flared LFs	2.24	1.15	0.57	0.50	0.44	0.40	0.35
LFGTE LFs	4.39	2.73	1.40	1.26	1.14	1.04	0.95
Industrial LFs	0.92	0.88	0.78	0.70	0.63	0.58	0.53
Out-of-State LFs	2.21	1.65	0.97	1.58	1.67	1.75	1.81
Municipal WW	0.45	0.47	0.52	0.54	0.57	0.61	0.64
Total Waste Management	13.38	9.61	6.67	6.76	6.46	6.21	6.01

Key Uncertainties

The methods used to model landfill gas emissions do not adequately account for the points in time when controls were applied at individual sites. Hence, for landfills, the historical emissions are less certain than current emissions and future emissions for this reason (since each site that is currently controlled was modeled as always being controlled, the historic emissions are low as a result). The modeling also does not account for uncontrolled sites that will need to apply controls during the period of analysis due to triggering requirements of the federal New Source Performance Standards/Emission Guidelines. As noted above, the available data do not cover all of the open and closed landfills in New Jersey. For this reason, emissions could be slightly underestimated for landfills.

For industrial landfills, these were estimated using national defaults (7% of the rate of MSW emplacement). It could be that the available MSW emplacement data within the NJDEP data used to model the MSW emissions already captures industrial landfill emplacement. As with overall MSW landfill emissions, industrial landfill emissions are projected to decrease between 2005 and 2020. If separate industrial landfill sites do not exist and the existing municipal waste emplacement data are thought to include industrial wastes, then the industrial landfill emissions can be excluded from the inventory.

Waste that is transported out of state was modeled assuming the same level of controls as the New Jersey landfills. Variances from this control level are possible, based on the mix of controls used by the receiving landfills. This analysis also assumed that the share of the State's waste disposed of out-of-state remains constant in the future. It should be noted that, some GHG inventories for other states receiving waste from New Jersey will include these emissions within that state's inventory as well⁹¹.

As noted, emissions from residential open burning (e.g. in backyard burn barrels) are not included. While the additional emissions are likely to be small, estimates can be made if open burning activity data are identified.

For the wastewater sector, the key uncertainties are associated with the application of SGIT default values for the parameters listed in Table G1 above (e.g., fraction of BOD which is anaerobically decomposed). The SGIT defaults were derived from national data.

As described above, data were not available to estimate emissions from the industrial wastewater treatment sector. If these data are identified (either production volumes in each industry or wastewater flow & chemical oxygen demand estimates), then emission estimates can be derived.

⁹¹ The emissions associated with NJ waste will only be included in another state's inventory if it uses historical data on total amounts of waste deposited in the state. If instead a state uses the SGIT default approach, its CH₄ emissions from landfills will be estimated using population and national per capita estimates of amount of waste landfilled, and thus will not capture the emissions associated with waste imported from NJ or elsewhere.

Appendix H. Forestry and Land-Use Change

Overview

Forestland emissions refer to the net carbon dioxide (CO₂) flux⁹² from forested lands in New Jersey, which account for about 40% of the state's land area.⁹³ The dominant forest type in New Jersey is Oak-Hickory which makes up about 46% of forested lands. Other common forest types are Loblolly-Shortleaf Pine (30%) and Oak-Gum-Cypress, Maple-Beech-Birch, and Oak-Pine, which make up less than 10% each.

Through photosynthesis, carbon dioxide is taken up by trees and plants and converted to carbon in biomass within the forests. Carbon dioxide emissions occur from respiration in live trees, decay of dead biomass, and fires. Carbon is also stored in the associated soils. In addition, carbon is stored for long time periods when forest biomass is harvested for use in durable wood products. Carbon dioxide flux is the net balance of carbon dioxide removals from and emissions to the atmosphere from the processes described above.

Emissions of carbon dioxide occur from the loss of forests and other vegetated lands when these are cleared for construction of buildings, parking lots, etc.⁹⁴ Estimates of these emissions of carbon dioxide can be combined with estimates of the net carbon dioxide flux from forested and other vegetated lands to develop a combined estimate of the overall net carbon dioxide flux associated with land-use and land-use change.

The following discusses the data sources, methods, assumptions, and results used to construct the inventory and reference case projections from forestry and land-use change. The reference case projections assume business-as-usual practices.

Emissions and Reference Case Projections (Business-as-Usual)

There are two methods that have been used as part of this project to estimate the net flux of carbon dioxide associated with forest growth.

Method 1

This approach makes use of the carbon online estimator (COLE),⁹⁵ and couples this with statewide estimates of different types of land cover. COLE was used by the New Jersey Department of Environmental Protection (NJDEP) to develop an estimate that, statewide, an average of 4.8 metric tons (Mt) of carbon dioxide equivalents (CO₂e) are sequestered per hectare per year in New Jersey from forested lands.⁹⁶

⁹² "Flux" refers to both emissions of CO₂ to the atmosphere and removal (sinks) of CO₂ from the atmosphere.

⁹³ Total forested acreage is 1.92 million acres in 2004; J. Smith, USFS, personal communication with S. Roe, CCS, April 2007. Acreage by forest type available from the USFS at: <http://www.fs.fed.us/ne/global/pubs/books/epa/states/NJ.htm>. The total land area in New Jersey is 4.75 million acres (<http://www.50states.com/newjerse.htm>).

⁹⁴ It is assumed herein that these emissions are associated with a non-sustainable loss of biomass, although in some cases, e.g. residential developments, trees are replanted and their growth will sequester carbon

⁹⁵ <http://ncasi.uml.edu/COLE/>

⁹⁶ Reyes, Jorge, NJDEP, 2007, personal communication

There are three different sources of estimates of the changes in land cover over time in the state. One of these is the Forest Inventory Analysis (FIA). The FIA data show a net increase in forest area in the 1987-1999 period of 243,000 acres and a decrease in area of 214,000 acres in the 1999-2004 period.⁹⁷ Another is the aerial photography completed by the State of New Jersey as interpreted by the NJ Department of Environmental Protection's Geographical Information Systems (GIS) unit . These data are shown in Table H1.

Table H1. New Jersey Land-Use Type, Acres, from NJDEP/GIS

Year	Developed	Crop/Grass	Upland Forest	Bare	Wetlands
1986	1,208,553	1,006,980	1,641,279	38,450	788,870
1995	1,334,565	883,590	1,616,525	45,530	737,010
2002	1,440,464	849,999	1,575,220	58,982	734,028

A third source of land cover data is the Center for Remote Sensing and Spatial Analysis, at Rutgers University (CRSSA).⁹⁸ These data are as in Table H2.

Table H2. New Jersey Land-Use Type, Acres, from CRSSA

Year	Developed	Cultivated & Grassland	Upland Forest	Bare	Inland Wetlands
1972	888,520	999,340	1,673,110	29,840	925,300
1984	1,204,920	1,006,980	1,465,680	38,450	788,870
1995	1,427,315	883,590	1,421,060	45,530	737,010
2001	1,483,158	849,999	1,388,941	58,982	734,028

With the assumption that inland wetlands, as shown in Table H2, are largely forested, the average rate of change of forested land based on all three data sets combined for the period from 1984 to 2002 was determined to be a loss of 1.2% of forested acreage per year.⁹⁹ This rate was used to project total forested acres into the future, which were then multiplied by the figure of 4.8 Mt CO₂e sequestered per hectare per year as determined by the COLE method as noted above to estimate the total statewide amount of CO₂e sequestered in any given year. These estimates are shown in Table H3.

⁹⁷ Smith, Jim, U.S. Forest Service, USFS, personal communications with S. Roe, CCS, October 2006 and May 2007.

⁹⁸ <http://www.crssa.rutgers.edu/projects/lc/>, accessed 6/21/07

⁹⁹ In this calculation, the FIA data were assumed to represent points on a linear trend; the linear function was estimated and acreages in intermediate years were determined by interpolation.

Table H3. Method 1 Estimated CO₂ Equivalent Sequestration, New Jersey

Year	Millions of Metric Tons CO ₂ Equivalent Sequestered per Year
1990	7.81
1995	7.37
2000	6.95
2005	6.55
2010	6.18
2015	5.83
2020	5.50

Releases of carbon from the loss of forests and other vegetated lands when these are cleared for construction of buildings, parking lots, etc. were estimated based on several data sources as approximately 95 Mt per acre.¹⁰⁰ When converted to CO₂e¹⁰¹, and multiplied by one half of the estimated increase in developed area per year determined from the NJDEP GIS and CRSSA data (with the assumption that half of the land converted to developed land each year was originally forested), a statewide total CO₂ equivalent release of approximately 1.1 million metric tons (MMt) per year throughout the 1984 to 2002 period was estimated.

Method 2

This approach relied on NJDEP/GIS land-use data (Table H1) and relevant default assumptions¹⁰² about the quantity of stored carbon for each land-use type, including associated soil carbon. It also took into consideration the carbon remaining in harvested wood products. Carbon remains stored in the durable wood products¹⁰³ pool or is transferred to landfills where much of the carbon remains stored over a long period of time.

With this method, the amount of land remaining in each land-use type and also the amount of land lost from each land type¹⁰⁴ was multiplied by factors based on the quantity assumptions listed above including annual biomass growth and soil density increase. Overall total net quantities of the equivalent CO₂ sequestered for the periods 1986 to 1995 and 1995 to 2002 were estimated to be 6.18 MMt tons per year for the first period and 5.93 MMt per year for the second period.

¹⁰⁰ These include the following: 1), the default value from the Intergovernmental Panel on Climate Change (IPCC) of 87.5 to 125 metric tons carbon/ha (based on a dry matter/ha estimate from this source, and with the assumption that the dry matter is 50% carbon); 2), calculations by Jorge Reyes, NJDEP which result in a figure of 96 metric tons carbon/ha; 3), Table A-191 from Annex 3 of the USEPA's 2007 Draft Greenhouse Gas Emissions inventory, which indicates that above-ground matter in the Northeastern U.S. forests contains about 103 metric tons carbon/ha, 4) an estimate by Ed Lempicki, NJDEP Division of Forestry, that there are 80 tons of biomass material per acre, which, , converts to 98 metric tons carbon/ha; and 5),

¹⁰¹ Based on the following conversion: $MTCO_2 = MTC \times 3.66$

¹⁰² Adapted from IPCC and other sources: [a] Biomass carbon density: 38 metric tons (Mt)/acre (forest), 4 Mt/acre (grassland), 2 Mt/acre (bare land), 1.2 Mt/acre (cropland); [b] soil carbon density: 8 Mt/acre (bare land) and 24 Mt/acre (forest land); [c] biomass density increase: 1% per year; [d] soil carbon density increase: 1% per year; and [e] amount of carbon stored in forest products: 12 Mt/acre.

¹⁰³ We estimate 50% of forest removal converted to wood products. Factor used to convert wood volume to weight: 3 pounds per board foot.

¹⁰⁴ In the absence of default factors for wetlands, we assumed them to be 50% forested and 50% bare.

Methods 1 and 2 Combined

A comparison of the estimates developed with Method 2 and net yearly sequestration estimates derived from taking the difference between sequestered and released quantities developed with Method 1 reveals close agreement. An average of both methods' net sequestration quantities for the 1986 to 1995 period is approximately 6.4 MMt per year and for the 1995 to 2002 period is approximately 6.0 MMt per year. With the assumption that the rate of decline suggested by these two average values is linear and continues at the same rate into the future, the statewide net sequestration from the landscape for 2004 is estimated as approximately 5.7 MMt, declining to 5.6 MMt in 2005, 5.4 MMt in 2010, 5.1 MMt in 2015, and 4.8 MMt in 2020. Table H4 shows the resulting sequestration values assumed for New Jersey. All values were calculated based on the linear rate of decline derived from the average values for the two time periods, and assuming that approximately 1.1 MMtCO₂e per year are released statewide through the clearing of forests and vegetated lands.

Table H4. Resultant Estimated CO₂ Equivalent Sequestration, New Jersey

Year	CO ₂ Released per Year (MMtCO ₂ e)	CO ₂ Sequestered per Year (MMtCO ₂ e)	Net CO ₂ Sequestered per Year (MMtCO ₂ e)
1990	1.10	7.53	6.43
1995	1.10	7.26	6.16
2000	1.10	6.99	5.89
2005	1.10	6.73	5.63
2010	1.10	6.46	5.36
2015	1.10	6.19	5.09
2020	1.10	5.93	4.83

Key Uncertainties

It should be noted that methane (CH₄) and nitrous oxide (N₂O) emissions from wildfires and prescribed burns have not been included in the estimates presented herein. In work that the Center for Climate Strategies (CCS) has completed for a number of western states, where wildfire activity is significant, emission estimates have tended to range from <1 to 3 MMtCO₂e/yr. We expect that the emissions from wildfires in New Jersey would be much lower than these levels and to not impact the estimated flux significantly.

Emissions of CH₄ from anaerobic forest soils (e.g., swamps, wetlands) have not been captured in this assessment. This is an area that should be investigated in the future, pending availability of data, to provide a more comprehensive picture of GHG emissions from forest soils.