



COVID-19


People with Certain Medical Conditions

Updated Feb. 10, 2023

As of January 26, 2023, EVUSHELD™ is not currently authorized for emergency use because it is unlikely to be active against the majority of SARS-CoV-2 variants circulating in the United States.

If you or your family member are at high risk for severe illness, wear a mask or respirator with greater protection in public indoor spaces if you are in an area with a high COVID-19 Community Level. Talk with your healthcare provider about wearing a mask in a medium [COVID-19 Community Level](#).

If you have [symptoms](#) consistent with COVID-19 and you are aged 50 years or older OR are at [high risk](#) of getting very sick, you may be eligible for [treatment](#). Contact your healthcare provider and start treatment within the first few days of symptoms. You can also visit a [Test to Treat](#) [location](#). Treatment can reduce your risk of hospitalization by more than 50% and also reduces the risk of death.



This information is intended for a general audience. Healthcare professionals should see [Underlying Medical Conditions Associated with Higher Risk for Severe COVID-19](#) for more detailed information.

What You Need To Know

- A person with any of the medical conditions listed below is more likely to get very sick with COVID-19. If you have one of these conditions, talk with your healthcare provider about how best to protect yourself from severe illness from COVID-19.
- The list below does not include all possible conditions that put you at higher risk of severe illness from COVID-19. If you have questions about a condition not included on this list, talk to your healthcare provider about how best to manage your condition and protect yourself from COVID-19.
- [Staying up to date with COVID-19 vaccines](#) and following [preventive measures for COVID-19](#) are important. This is especially important if you are older or have severe health conditions or more than one health condition, including those on the list below.
- Approved and authorized COVID-19 vaccines (primary series and booster) are [safe and effective](#).
- [Some people who are immunocompromised](#), or people with weakened immune systems, may be eligible for an additional primary dose of COVID-19 vaccine.

Overview

Based on the current evidence, a person with any of the conditions listed below is more likely to get very sick from COVID-19. This means that a person with one or more of these conditions who gets very sick from COVID-19 (has severe illness from COVID-19) is more likely to:

- Be hospitalized
- Need intensive care
- Require a ventilator to help them breathe
- Die

In addition:

- [Older adults](#) are at highest risk of getting very sick from COVID-19. More than 81% of COVID-19 deaths occur in people over age 65. [The number of deaths among people over age 65 is 97 times higher than the number of deaths among people ages 18-29 years.](#)
- A person's risk of severe illness from COVID-19 increases as the number of underlying medical conditions they have increases.
- Some people are at increased risk of getting very sick or dying from COVID-19 because of where they live or work, or because they can't get health care. This includes many [people from racial and ethnic minority groups](#) and [people with disabilities](#).
 - Studies have shown people from racial and ethnic minority groups are also dying from COVID-19 at younger ages. People in racial and ethnic minority groups are often younger when they develop chronic medical conditions and may be more likely to have more than one medical condition.
 - People with disabilities are more likely than those without disabilities to have [chronic health conditions, live in shared group \(also called "congregate"\) settings, and face more barriers in accessing health care](#). Studies have shown that some people with certain disabilities are more likely to get COVID-19 and have worse outcomes.

[Staying up to date with COVID-19 vaccines](#) and [taking COVID-19 prevention actions](#) are important. This is especially important if you are older or have severe health conditions or more than one health condition, including those on this list. Learn more about how CDC develops [COVID-19 vaccination](#) recommendations. If you have a medical condition, learn more about [Actions You Can Take](#).

Medical Conditions

- The conditions on this list are in alphabetical order. They are **not in order of risk**.
- CDC completed a review for each medical condition on this list. This was done to ensure that these conditions met criteria for inclusion on this list. CDC conducts ongoing reviews of additional underlying conditions. If other medical conditions have enough evidence, they might be added to the list.
- Because we are learning more about COVID-19 every day, this list **does not** include all medical conditions that place a person at higher risk of severe illness from COVID-19. Rare medical conditions, including many conditions that mostly affect children, may not be included on the list below. We will update the list as we learn more.
- A person with a condition that is not listed may still be at greater risk of getting very sick from COVID-19 than other people who do not have the condition. It is important that you **talk with your healthcare provider about your risk**.

Cancer

Having cancer can make you more likely to get very sick from COVID-19. Treatments for many types of cancer can weaken your body's ability to fight off disease. At this time, based on available studies, having a history of cancer may increase your risk.


Get more information:

- [Cancer](#)
- [COVID-19: What People with Cancer Should Know – National Cancer Institute](#) [↗](#)

Chronic kidney disease

Having chronic kidney disease of any stage can make you more likely to get very sick from COVID-19.

Get more information:

- [Chronic kidney disease](#)
- [National Kidney Foundation: Kidney disease and COVID-19](#) 

Chronic liver disease

Having chronic liver disease can make you more likely to get very sick from COVID-19. Chronic liver disease can include alcohol-related liver disease, non-alcoholic fatty liver disease, autoimmune hepatitis, and cirrhosis (or scarring of the liver).

Get more information:

- [Liver Disease](#) 
- [American Liver Foundation: Your Liver & COVID-19](#) 

Chronic lung diseases

Having a chronic lung disease can make you more likely to get very sick from COVID-19. Chronic lung diseases can include:

- Asthma, if it's moderate to severe
- Bronchiectasis (thickening of the lungs' airways)
- Bronchopulmonary dysplasia (chronic lung disease affecting newborns)
- Chronic obstructive pulmonary disease (COPD), including emphysema and chronic bronchitis
- Having damaged or scarred lung tissue known as interstitial lung disease (including idiopathic pulmonary fibrosis)
- Pulmonary embolism (blood clot in the lungs)
- Pulmonary hypertension (high blood pressure in the lungs)

Get more information:

- [COPD](#)
- [Asthma](#)
- [People with Moderate to Severe Asthma](#)
- [American Lung Association: Controlling Chronic Lung Diseases Amid COVID-19](#) 

Cystic fibrosis

Having cystic fibrosis, with or without lung or other solid organ transplant (like kidney, liver, intestines, heart, and pancreas) can make you more likely to get very sick from COVID-19.

Get more information:

- [Cystic fibrosis](#)
- [CF and Coronavirus \(COVID-19\) | Cystic Fibrosis Foundation \(cff.org\)](#) 

Dementia or other neurological conditions

Having neurological conditions, such as dementia, can make you more likely to get very sick from COVID-19.

Get more information:

- [Dementia](#)
- [Alzheimer's Association: COVID-19, Alzheimer's and Dementia](#) 

Diabetes (type 1 or type 2)


Having either type 1 or type 2 diabetes can make you more likely to get very sick from COVID-19.

Get more information:

- [Diabetes](#)
- [American Diabetes Association: How COVID-19 Impacts People with Diabetes](#) 

Disabilities

People with some types of disabilities may be more likely to get very sick from COVID-19 because of underlying medical conditions, living in congregate settings, or systemic health and social inequities, including:

- [People with any type of disability that makes it more difficult to do certain activities or interact with the world around them, including people who need help with self-care or daily activities](#)
- [People with attention-deficit/hyperactivity disorder \(ADHD\)](#)
- [People with cerebral palsy](#)
- [People with birth defects](#)
- [People with intellectual and developmental disabilities](#)
- [People with learning disabilities](#)
- [People with spinal cord injuries](#) 
- [People with Down syndrome](#)

Get more information:

- [People with Disabilities | COVID-19](#)

Heart conditions

Having heart conditions such as heart failure, coronary artery disease, cardiomyopathies, and possibly high blood pressure (hypertension) can make you more likely to get very sick from COVID-19.

Get more information:

- [Heart Disease](#)
- [American Heart Association: COVID-19](#) 
- [NHLBI Information & Resources on COVID-19](#) 

HIV infection

Having HIV (Human Immunodeficiency Virus) can make you more likely to get very sick from COVID-19.

Get more information:

- [HIV Infection](#)
- [Interim Guidance for COVID-19 and Persons with HIV](#) 




Immunocompromised condition or weakened immune system

Some people are immunocompromised or have a weakened immune system because of a medical condition or a treatment for a condition. This includes people who have cancer and are on chemotherapy, or who have had a solid organ transplant, like a kidney transplant or heart transplant, and are taking medication to keep their transplant. Other people have to use certain types of medicines for a long time, like corticosteroids, that weaken their immune system. Such long-term uses can lead to [secondary or acquired immunodeficiency](#). Other people have a weakened immune system because of a life-long condition. For example, some people inherit problems with their immune system. One example is called [Primary immunodeficiency](#). Being immunocompromised can make you more likely to get very sick from COVID-19 or be sick for a longer period of time.

People who are immunocompromised or are taking medicines that weaken their immune system may not be protected even if they are [up to date on their vaccines](#). Talk with your healthcare provider about wearing a mask in a medium [COVID-19 Community Level](#) and what additional precautions may be necessary in medium or high COVID-19 Community Levels.

After completing the primary vaccination series, some people who are [moderately or severely immunocompromised](#) should get an additional primary dose and a booster. Because the immune response following COVID-19 vaccination may differ in people who are moderately or severely immunocompromised, [specific guidance](#) has been developed.




Get more information:

- [Types of Primary Immune Deficiency Diseases](#) 
- [Jeffrey Modell Foundation](#) 
- [Immune Deficiency Foundation](#) 
- [Primary Immunodeficiency \(PI\)](#)

Mental health conditions

Having mood disorders, including depression, and schizophrenia spectrum disorders can make you more likely to get very sick from COVID-19.


Get more information:

- [National Institute of Mental Health \(NIMH\) Shareable Resources on Coping with COVID-19](#) 
- [National Institute of Mental Health \(NIMH\) Depression](#) 
- [Mood Disorders](#) 

Overweight and obesity




Overweight (defined as a [body mass index](#) (BMI) is 25 kg/m² or higher, but under 30 kg/m²), obesity (BMI is 30 kg/m² or higher, but under 40 kg/m²), or severe obesity (BMI is 40 kg/m² or higher), can make you more likely to get very sick from COVID-19. The risk of severe illness from COVID-19 increases sharply with higher BMI.

Get more information:

- [Overweight & Obesity](#)
- [Obesity, Race/Ethnicity, and COVID-19](#)
- [Obesity Action Coalition: COVID-19 and Obesity](#) 

Physical inactivity

People who do little or no physical activity are more likely to get very sick from COVID-19 than those who are physically active. Being physically active is important to being healthy. Get more information on physical activity and health, physical activity recommendations, how to become more active, and how to create activity-friendly communities:

- [Physical Activity](#)
- [Physical Activity Guidelines for Americans, 2nd edition](#) 
- [Move Your Way®](#) 
- [Active People, Healthy Nation SM: Strategies to Increase Physical Activity](#)
- [National Center on Health, Physical Activity and Disability – Building Healthy Inclusive Communities](#) 

Pregnancy

Pregnant and recently pregnant people (for at least 42 days following end of pregnancy) are more likely to get very sick from COVID-19 compared with non-pregnant people.

Get more information:

- [Pregnant and Recently Pregnant People](#)
- [Toolkit for Pregnant People and New Parents](#)

- [Investigating the Impact of COVID-19 during Pregnancy](#)

Sickle cell disease or thalassemia

Having hemoglobin blood disorders like sickle cell disease or thalassemia (inherited red blood cell disorders) can make you more likely to get very sick from COVID-19.

Get more information:

- [Sickle Cell Disease](#)
- [Thalassemia](#)

Smoking, current or former

Being a current or former cigarette smoker can make you more likely to get very sick from COVID-19. If you currently smoke, quit. If you used to smoke, don't start again. If you've never smoked, don't start.

Get more information:

- [Smoking & Tobacco Use](#)
- [Tips From Former Smokers](#)
- [Health Benefits of Quitting Smoking](#)

Solid organ or blood stem cell transplant

Having had a solid organ or blood stem cell transplant, which includes bone marrow transplants, can make you more likely to get very sick from COVID-19.

Get more information:

- [Transplant Safety](#)
- [COVID-19 Resources for Transplant Community](#) 

Stroke or cerebrovascular disease

Having cerebrovascular disease, such as having a stroke which affects blood flow to the brain, can make you more likely to get very sick from COVID-19.


Get more information:

- [Stroke](#)
- [COVID19 Stroke Podcast Series for Patients and Caregivers](#) 

Substance use disorders

Having a substance use disorder (such as alcohol, opioid, or cocaine use disorder) can make you more likely to get very sick from COVID-19.

Get more information:

- [How to Recognize a Substance Use Disorder](#) 
- [Drug Overdose](#)

Tuberculosis

Having tuberculosis (TB) can make you more likely to get very sick from COVID-19.

Get more information:

- [Basic TB Facts](#)
- [Public Health Emergencies](#)

Additional Information on Children and Teens

People of all ages, including children, can get very sick from COVID-19. Children with underlying medical conditions are at increased risk for getting very sick compared to children without underlying medical conditions.

Current evidence suggests that children with medical complexity, with genetic, neurologic, or metabolic conditions, or with congenital heart disease can be at increased risk for getting very sick from COVID-19. Like adults, children with obesity, diabetes, asthma or chronic lung disease, sickle cell disease, or who are immunocompromised can also be at increased risk for getting very sick from COVID-19. Check out [COVID-19 Vaccines for Children and Teens](#) for more information on vaccination information for children.

- [COVID-19 Vaccines for Children and Teens](#)

Actions You Can Take

It is important to protect yourself and others by taking [COVID-19 prevention actions](#):

- Stay up to date with your COVID-19 vaccines
- Improve ventilation
- Get tested if you have symptoms
- Follow recommendations for what to do if you have been exposed
- Stay home if you have suspected or confirmed COVID-19
- Seek treatment if you have COVID-19 and are at high risk of getting very sick
- Avoid contact with people who have suspected or confirmed COVID-19
- Wear a mask or respirator
- Increase space and distance

Please [contact your state, tribal, local, or territorial health department](#) for more information on [COVID-19 vaccination in your area](#). It is also important for people with medical conditions and their healthcare providers to work together and manage those conditions carefully and safely. [Stay up to date with your COVID-19 vaccines](#). If you have a medical condition, the following are actions you can take based on your medical conditions and other risk factors.

Build Your Personal COVID-19 Plan

COVID-19 Plan
Tools, information, and action steps to share with your family, friends, and healthcare provider

Start your personal COVID-19 plan
Talk with your healthcare provider about whether you are at high-risk of getting very sick from COVID-19.

- People who are more likely to get very sick include older adults (ages 50 years or more, with risk increasing with age), people who are unvaccinated, and people with certain medical conditions, such as chronic lung disease, heart disease, or a weakened immune system.

Understand risk
<https://www.cdc.gov/coronavirus/2019-nCoV/your-health/understanding-risk.html>

- People with risk factors for severe disease may benefit from treatment if they get COVID-19. A healthcare provider will help decide which treatment, if any, is right for you.

Tests
Have a supply of COVID-19 self-test kits at home and know when and how to use them.
<https://www.cdc.gov/coronavirus/2019-nCoV/testing/self-testing.html>

- If you have symptoms of COVID-19, test immediately.
- If you test positive, treatments are available that can reduce your chances of hospitalization and death.

Treatment
Have a plan to contact a healthcare provider right away if you test positive.

- Don't delay: Treatment must be started within days after you first develop symptoms to be effective.

Healthcare provider
If you don't have a healthcare provider, consider telehealth options or contact a Test to Treat site to get tested, evaluated, and treated in one location.

- Test to Treat locations: <https://www.2019-test-to-treat.locator.dhs.gov/locations/>

Take precautions
COVID-19 is in your Community: Keep track of your COVID-19 Community Level and use it to guide your precautions. <https://www.cdc.gov/coronavirus/2019-nCoV/your-health/level-by-county.html>

Vaccines
Stay up to date on vaccines. Know when to get a booster.
<https://www.cdc.gov/coronavirus/2019-nCoV/your-health/stay-up-to-date.html>

Ventilation
Spend time outside and improve indoor air quality at home by opening windows and using other strategies.

Masks
Masks are recommended for those at high risk when COVID-19 community levels are medium and for everyone when levels are high. Learn more: <https://www.cdc.gov/coronavirus/2019-nCoV/prevent-getting-sick/about-face-coverings.html>

Put together your COVID-19 plan so you have all the information you need on hand if you get sick with COVID-19. Download, edit and save, and share your plan with your family, friends, and healthcare provider.

File Details: 561 KB, 3 pages

[View PDF in English](#)

[Other Languages](#)

Seek care when needed

- Call your healthcare provider if you have any concerns about your medical conditions or if you get sick and think that you may have COVID-19. Discuss steps you can take to manage your health and risks. **If you need emergency help, call 911 right away.**
- Do not delay getting care for your medical condition because of COVID-19. Emergency departments, urgent care, clinics, and your healthcare provider have infection prevention plans to help protect you from getting COVID-19 if you need care.

Continue medications and preventive care

- Continue your medicines and do not change your treatment plan without talking to your healthcare provider.
- Have at least a 30-day supply of prescription and non-prescription medicines. [Talk to a healthcare provider](#), insurer, or pharmacist about getting an extra supply (i.e., more than 30 days) of prescription medicines, if possible, to reduce your trips to the pharmacy.
- Follow your current treatment plan (e.g., [Asthma Action Plan](#), dialysis schedule, blood sugar testing, nutrition, and exercise recommendations) to keep your medical condition(s) under control.
- When possible, keep your appointments (e.g., vaccinations and blood pressure checks) with your healthcare provider. Check with your healthcare provider about safety precautions for office visits and ask about telemedicine or virtual healthcare appointment options.
- Learn about [stress and coping](#). You may feel increased stress during this pandemic. Fear and anxiety can be overwhelming and cause strong emotions. It can be helpful to talk with a professional like a counselor, therapist, psychologist, or psychiatrist. Ask your primary care provider if you would like to speak with a professional. Getting regular exercise and being physically active is also a great way to reduce stress.

Accommodate dietary needs and avoid triggers

- Have non-perishable food choices such as canned goods available that meet your needs based on your medical condition (e.g., kidney diet and [KCER 3-Day Emergency Diet Plan](#) [↗](#), diabetic diet).
- Know the triggers for your condition and avoid when possible (e.g., avoid [asthma triggers](#) by having another member of your household clean and disinfect your house for you or [avoid possible sickle cell disease triggers](#) to prevent pain crises).

Last Updated Feb. 10, 2023